



Review

Inflammation, Dopaminergic Brain and Bilirubin

Sri Jayanti ^{1,2} , Camilla Dalla Verde ¹ , Claudio Tiribelli ^{1,*} and Silvia Gazzin ¹

¹ Italian Liver Foundation, Liver Brain Unit “Rita Moretti”, Area Science Park, Bldg. Q, SS 14, Km 163,5, 34149 Trieste, Italy; sri.jayanti@fegato.it (S.J.); camilla.dallaverde@fegato.it (C.D.V.); silvia.gazzin@fegato.it (S.G.)

² Eijkman Research Centre for Molecular Biology, Research Organization for Health, National Research and Innovation Agency, Cibinong 16915, Indonesia

* Correspondence: ctliver@fegato.it

Abstract: Dopamine is a well-known neurotransmitter due to its involvement in Parkinson’s disease (PD). Dopamine is not only involved in PD but also controls multiple mental and physical activities, such as the pleasure of food, friends and loved ones, music, art, mood, cognition, motivation, fear, affective disorders, addiction, attention deficit disorder, depression, and schizophrenia. Dopaminergic neurons (DOPAn) are susceptible to stressors, and inflammation is a recognized risk for neuronal malfunctioning and cell death in major neurodegenerative diseases. Less is known for non-neurodegenerative conditions. Among the endogenous defenses, bilirubin, a heme metabolite, has been shown to possess important anti-inflammatory activity and, most importantly, to prevent DOPAn demise in an ex vivo model of PD by acting on the tumor necrosis factor-alpha (TNF α). This review summarizes the evidence linking DOPAn, inflammation (when possible, specifically TNF α), and bilirubin as an anti-inflammatory in order to understand what is known, the gaps that need filling, and the hypotheses of anti-inflammatory strategies to preserve dopamine homeostasis with bilirubin included.

Keywords: addiction; cognition; depression; schizophrenia; Parkinson’s disease; attention deficit hyperactivity disorder; tumor necrosis factor; PPARs; NF κ B; MAPKs



Citation: Jayanti, S.; Dalla Verde, C.; Tiribelli, C.; Gazzin, S. Inflammation, Dopaminergic Brain and Bilirubin. *Int. J. Mol. Sci.* **2023**, *24*, 11478. <https://doi.org/10.3390/ijms241411478>

Academic Editor: Yingfang Tian

Received: 28 June 2023

Revised: 7 July 2023

Accepted: 11 July 2023

Published: 14 July 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The impairment of the physiologic dopamine (DA) function and the death of dopaminergic neurons (DOPAn) bring Parkinson’s disease immediately to mind; however, DA affects extensive tasks in the brain. DA is involved in pleasure, sleep, mood, cognition, motivational salience, reward-motivated behavior, hormone release, addiction, depression, schizophrenia, attention deficit hyperactivity disorder, multiple sclerosis, ischemic stroke, and more [1–6]. Thus, the homeostasis of DA systems exceeds what would generally be perceived, involving both mental and physical health.

Like adrenaline and noradrenaline, DA is a catecholamine that possesses the 3,4-dihydroxy aromatic ring (the catechol core) synthesized from L-tyrosine, which could be detected in the whole body. The DA produced in the brain cannot pass the blood–brain barrier (BBB); however, peripheral organs can produce it and react with it thanks to the presence of five different receptors, which, in turn, act on multiple signaling pathways (see later). The DA cascade might also finally play a critical role in mammals’ whole body physiology, health, and diseases [7]. For example, DA controls the pathophysiology of the heart and cardiovascular system, gastrointestinal system, liver, innate and adaptive immunity, the kidney (see also Section 4), and the brain [7–11].

In the brain, DA is synthesized through a restricted population of cells (less than 1% of the total number of neurons in the central nervous system (CNS) that are located in specific areas [1,12,13] and form four major dopaminergic circuits (see later and Figure 1)). Of importance, DA could regulate both innate and adaptive immunity due to its ability to

bind to cell receptors of both immune and neuronal systems as well those reviewed by Carandini et al. [14] and Mingan Li [11]. On the other hand, systemic inflammation has been reported to affect a broad range of neurologic and neurodegenerative conditions, with aging leading to a 4–7% DOPAn loss per decade and environment and lifestyle affecting the systemic inflammatory status and amplifying the mechanisms [10,11,14–22].




Figure 1. Cartoon of the four dopaminergic circuits, their major function, and the pathologic conditions affecting each circuit. Red: the mesocortical circuit; black: mesolimbic circuit; blue: nigrostriatal circuit; and green: tubero-infundibular circuit. Am: amygdala; Ac: nucleus accumbens; Cc: corpus callosum; Cn: caudate nucleus; fCtX: frontal cortex; Ot: optic tract; Pg: pituitary gland; R: retina; VTA: ventral tegmental area; Zi: zona incerta. Circles with numbers at the origin of each circuit indicate the histologic classification of the dopaminergic neurons. PD: Parkinson's disease, HD: Huntington's disease, ADHD: attention deficit hyperactivity disorder.

The brain and the body have increasingly been suggested as a single entity in terms of inflammation. Neuroinflammation is considered a driving mechanism and a possible marker in the occurrence, diagnosis, and treatment of neurodegenerative diseases and neurologic conditions [23]. In the case of DA, studies have consistently reported evidence of

a correlation between increased chronic low-grade inflammation, elevated peripheral and central inflammatory cytokines, inflammatory mediators, and acute-phase reactants (interleukin: IL6, IL1; tumor necrosis factor: TNF α ; C-reactive protein—CRP, etc.), with decreased DA availability as a subject with neuropsychiatric disorders [14,22,24]. Indeed, infectious diseases (HIV, sepsis), trauma, and TNF α administration as therapeutics in cancer have also been reported to induce depressive symptoms, with reduced motivation, anhedonia, and altered sleep [24]. Signaling pathways connecting inflammation and DOPAn degeneration have been suggested and evaluated as potential therapeutic targets. Among the most reported include cAMP/PKA/MAPK (cyclic adenosine monophosphate/protein kinase A/mitogen-activated protein kinases), NRF2 (nuclear factor erythroid 2-related factor 2), PPARs (peroxisome proliferator-activated receptors), NF κ B (nuclear factor kappa-light-chain-enhancer of activated B cells), TLR4 (toll-like receptor), and inflammasome-NLRP3 (NOD-, LRR- and pyrin domain-containing protein 3) [4,10,11,15,18,22,25–28] (Figure 2).




Figure 2. Dopamine and bilirubin metabolism, biological effects, and shared signaling pathway. **(a)** Cartoon showing dopamine (left) and bilirubin (right) metabolism. **(b)** Cartoon resuming the principal

effectors of DA dysregulation in neuropsychiatric and neurodegenerative conditions described in this paper (left), and the potential beneficial effects of YPS based both on clinic and experimental information (right), and the main signaling pathway shared between the neurologic conditions and YPs (middle). ↑: increase; ↓: decrease; TH: tyrosine hydroxylase; L-DOPA: L-dihydroxyphenylalanine; DA: dopamine; MAO: monoamine oxidase; DOPAL: 3,4-dihydroxyphenylacetaldehyde; ALDH: aldehyde dehydrogenases; DOPAC: 3,4-dihydroxyphenylacetic acid; COMT: catechol-o-methyltransferase; HVA: homovanilic acid. Heme: hemoglobin; HMOX: heme oxygenase; BV: biliverdin; BLVR: biliverdin reductase; UCB: unconjugated bilirubin («bilirubin»); UGT1a1: uridine-glucuronosyl transferase 1a1; CB: conjugated bilirubin; ROS: reactive oxygen species; noOS: unreactive oxygen species; PD: Parkinson's disease; NOS: nitrogen reactive species; COX2: cyclooxygenase 2; CXCL12: C-X-C motif chemokine ligand 12; TNF: tumor necrosis factor; IL: interleukin; TLR4: *Toll-Like Receptor 4*; MAPK: mitogen-activated protein kinase; NFκB: Nuclear factor NF-kappa-B; RE: responsive elements; TNF-R1: tumor necrosis factor receptor 1; ADHD: attention deficit hyperactivity disorder; ASD: autism spectrum disorder; IFN: interferon; Dp: depression; PPAR: peroxisome proliferator-activated receptors; Add: addiction; DR5: dopamine receptor 5; MS: multiple sclerosis; Sch: schizophrenia; EAE: experimental autoimmune encephalomyelitis (model of MS. multiple sclerosis).

Bilirubin is another body metabolite with pleiotropic and surprising functions, both at systemic and CNS levels. Mainly known as a waste product of hemoglobin catabolism possibly induces neuronal damage in infants and as a marker of liver dysfunction over the last decades, bilirubin has been rediscovered as a homeostatic, anti-oxidant, anti-inflammatory factor, and a hormone [29–35]. Epidemiologic studies have repeatedly reported an inverse correlation (protective) between mildly elevated total serum bilirubin (TSB) and chronic inflammatory conditions (e.g., diabetes, metabolic syndrome, cardiovascular diseases). At the same time, a low TSB (about or below the lower physiologic range of 0.4–1 mg/dL) has been associated with neurological conditions, suggesting that this pigment is a risk factor or a marker of CNS diseases [30,31,35]. The therapeutic application of bilirubin has started to be discussed [29,36,37]. Understanding the possible biomolecular effects on different neurologic diseases is mandatory from this perspective. Bilirubin is produced by the action of two sequential reactions. The first one is the conversion of hemoglobin to biliverdin, which is played by heme oxygenase (HMOX1); subsequently, biliverdin (BV) is converted to unconjugated bilirubin (UCB, or indirect bilirubin, simply bilirubin further in the review) by biliverdin reductase (BLVR) (Figure 2). Bilirubin, BV, HMOX1, and BLVR (the yellow players -YPs) activity and modulation in the course of this disease have been documented in the brain [35,38–43]. This high capability to modulate their level and activity, joined with documented benefits, has supported the idea that YPs are a form of homeostatic and defensive machinery in the cell. Moreover, YPs can act directly and indirectly (through signaling pathways, including—PPAR; MAPK; NRF2; AhR—aryl hydrocarbon receptor; NFκB; insulin pathway IRs/IRK/PI3K—insulin receptor substrates/insulin transmembrane receptor kinase/phosphatidylinositol 3-kinase [30,31,35]) on inflammation (e.g., TNFα, IL6, complement, T cell response, preventing the alteration of the BBB [44]). On the other hand, a sustained pro-inflammatory status usually down-regulates the YPs [44]. YPs are primarily known to modulate the cellular redox balance: another mechanism that is often reported in neurologic disease. According to the available data, this review aims to discuss the evidence connecting DOPAn, inflammation, and bilirubin.

2. The Dopamine Circuits in Brain

An in-depth review of the dopamine pathway is out of the scope of this review. The following section aims to provide a laid background on the DA pathway and DOPAn location in the brain to help readers visualize the complexity of dopamine's function in human behavior and related neurologic conditions.

To accomplish this objective, we performed a search of the literature on PubMed, Scopus, and Google Scholar as follows. (1) First, we searched for the most focused and relevant works in the literature using the keywords “dopamine”, or “dopamine system”, or

“dopamine diseases/pathology and inflammation”. (2) Second, we searched for “dopamine and bilirubin, or HMOX, or BLVR, or BV”. To this step, we added the literature that was already part of our background on YPs. (3) Finally, after reading the most relevant publications, we refined the literature search using keywords that were the most promising specific terms to emerge from steps 1 and 2.

DOPAn are a heterogenic family of nine different subtypes of cells (A8–14, A16, A17), which are grouped in well-defined brain regions connecting topographically organized anatomical tracts [1,3,12] that are mostly localized in mesencephalon and diencephalon (Figure 1). They all express genes that are critical for dopamine catabolism (Figure 2) and survival [6,12,45].

DA is involved in four major circuits in the central nervous system: (a) The nigrostriatal circuitry; (b) The tuberoinfundibular circuitry; (c) The mesocortical circuitry; and (d) the mesolimbic circuitry, with the last two largely overlapped, thus possibly referring together to mesocorticolimbic circuitry [46]. Altogether the four DA circuits described accounted for 90% of DOPAn in the brain [1]. Additional DOPAn were located in the retina, olfactory tubercle, and zona incerta [12] (Figure 1). The biology of DA in inflammatory conditions might be impressively variable, complicating the understanding of therapeutic approaches. These effects could depend on DA synthesis, trafficking, sequestration into vesicles, and release, from the type of DA receptors expressed in target tissues and cells (DA receptors: DR, G-coupled receptors type; with DR2, DR3, and DR4 usually inhibitory; and DR1, DR5 usually excitatory) from the concentration of the cytokine, the duration of the stimuli and even by the DA circuit involved [4,11,19,22,47]. Finally, the dysregulation of DA activity resulted in pathophysiology [13,26,48–51].

2.1. The Nigrostriatal Circuitry

The nigrostriatal circuit originates from a small group of DOPAn (histologic identification A8, 9; 3–5% of total neurons of the area) located in the substantia nigra pars compacta (SNpc) and projects to the basal ganglia (dorsal striatum: caudate nucleus and putamen). This pathway mainly regulates voluntary body movement but has also been involved in procedural and associative learning, including speech, cognition, reward, and addiction. The neuropsychiatric and neurodegenerative conditions, usually referred to as the nigrostriatal circuit, accounted for HD, attention-deficit hyperactivity disorder, schizophrenia, bipolar disorders, and obviously, Parkinson’s disease (PD) [12,13,50,52–60].

2.2. The Mesolimbic Circuitry

The DOPAn populating the ventral tegmental area (VTA, A10, 60–65% of the neurons of this region) projects to the ventral striatum (nucleus accumbens—Na, olfactory tubercle—Ot), the amygdala—Am, the hippocampus, and septum. This circuit is mainly involved in all instinctual and emotion-based behaviors (Figure 1), such as motivation, aversion, incentive salience, pleasure, reinforcements, reward, the determination of personality traits, and cognition. The dysregulation of this mesolimbic circuit underlies a whole plethora of neuropsychiatric diseases, including among them addiction, attention-deficit hyperactivity disorder, schizophrenia, depression, and obsessive-compulsive behaviors, as well as chronic pain. If usually the perturbation of the mesolimbic DA reward system has been linked to addiction to drugs, the correct meaning of the term is worth mentioning in that the same pathway could create a good lifestyle, such as “addiction” to sports, good food, responsible behavior for drugs, alcohol, etc. It may be of high relevance in adolescents [12,13,19,48,61–64].

2.3. The Mesocortical Circuitry

The DOPAn of the VTA (A10) also projects to the prefrontal cerebral cortex, the cingulate cortex, and the perirhinal cortex (Figure 1). This circuit is implicated in cognition, executive functions, emotions, learning and memory, pleasure, and chronic pain. Depression and schizophrenia are frequently related to the malfunctioning of this DA circuit [12,13,62–68].

2.4. The Tuberoinfundibular Circuitry

In the tuberoinfundibular circuit, DOPAn secretes DA from the arcuate (A12) and periventricular nuclei of the hypothalamus (A14) to the anterior pituitary gland. This pathway controls the hypothalamic-pituitary endocrine system and inhibits prolactin secretion from the anterior pituitary gland into the blood, mainly affecting fertility and maternal behavior. As a matter of notice, treatment with anti-dopaminergic anti-psychotic drugs (as an example in schizophrenia) causes hyperprolactinemia (Figure 1) [12,13,50,69–71].

These DA circuits are interrelated, which could explain the presence of mixed cognitive, executive, behavioral, psychotic, and motor abnormalities in PD and other conditions involving dopamine [72].

3. Neurological Conditions Related to Dopamine

3.1. Parkinson's Disease

Parkinson's disease (PD) is the most known neurodegenerative condition and is characterized by a progressive reduction in DA and DOPAn, with the number of people affected predicted to double from 6.9 million in 2015 to 14.2 million in 2040 [73]. The loss of DA in SNpc is responsible for motor disabilities (mainly: bradykinesia, tremors, dystonia, gait complaints, and falls). Other non-motor symptoms (e.g., olfactory impairment and constipation, apathy, sleep abnormalities, anxiety, depression, together with cognition problems) are attributable to alterations in the mesocorticolimbic system that usually precede the motor symptoms [5,21,53,74,75]. The etiology of PD has not been fully elucidated. In 90% of cases, no genetic identifiable explanation can be obtained. Multiple possible triggers have been suggested, as reviewed by several authors. Among these triggers, environmental factors (metals, pesticides, and herbicides), lifestyle (especially the high consumption of milk and dairy products [76,77]), gut dysbiosis (by both the induction of a systemic pro-inflammatory status as well as by direct signaling through the vagus [21,78]), abuse substances (e.g., methamphetamine—by impairing DA transport and increasing its extracellular concentration), the metabolic syndrome (namely insulin resistance through PPARs signaling), brain trauma (by breaking the BBB and inducing inflammation), and physiologic aging, accompanied by the so-called “senescence neuroinflammation”, contribute or even are supposed to trigger DA loss [10,15,76,78].

Extensive clinical and experimental evidence supports the involvement of inflammation in the onset and progression of PD. Microgliosis, astrogliosis, and the CD8+ and CD4+ adaptive immune T cells present in the brain of PD specimens (human and models) can lead to the induction of inducible nitric oxide synthase (NOS) and cyclooxygenase (COX) alongside the release of the pro-inflammatory cytokine C-X-C motif chemokine ligand 12 (CXCL12), TNF α , interferon- γ (IFN γ), IL6 and IL1 β [44,79–81]. Specifically, TNF α has been suggested to be a relevant player in PD. TNF α elevation has been reported in both postmortem human PD specimens and models. It has been associated with an enhanced DOPAn demise, physical and cognitive decline severity, and correlated with the Hoehn and Yahr score [82–87] (Figure 3). Recently, an SNP (single nucleotide polymorphism) in the TNF α promoter gene (-1031C) has been associated with a rare form of early-onset idiopathic PD [88]), and TNF α (G308A) polymorphism has been associated with vascular Parkinson's complicated with pulmonary infection [89]. Interestingly, in the young onset forms of PD, symptoms frequently develop secondary to infectious diseases, stressing even more the association between the activation of the inflammatory cascade and DA neurodegeneration. Between 5 and 10% of PD cases may have a genetic cause. Again, interestingly, one of the most recent perspectives on the etiology of this disease has suggested that the genes involved in PD may share biochemical mechanisms or signaling pathways with inflammatory pathways [28,90–93].




Figure 3. The alteration in dopamine (light blue), inflammation factors (pink), and YPs (yellow) in DA-related diseases. ↑: increase; ↓: decrease; DOPAn: dopaminergic neuron; DA: dopamine; CXCL12: C-X-C Motif Chemokine Ligand 12, TNF α : tumor necrosis factor α , IFN γ : interferon γ , IL6: interleukin 6, IL1 β : interleukin 1 β ; HMOX1: heme oxygenase 1; TSB: total serum bilirubin; DR: dopamine receptor; DR5: dopamine receptor 5; COX2: cyclooxygenase 2; IFN: interferon; IL1: interleukin 1, UCB: unconjugated bilirubin, DAT: dopamine transporter; DR1: dopamine receptor 1, DR3: dopamine receptor 3, DR4: dopamine receptor 4, IL2: interleukin 2; GABA: gamma-aminobutyric acid; Na: nucleus accumbens, Am: amygdala; IL8: interleukin 8, THF: of 6,7,4'-trihydroxyflavone. ADHD: Attention Deficit Hyperactivity Disorder; ASD: autism spectrum disorder.

Levodopa, DA agonists, MAO (monoamine oxidase) inhibitors, and other non-pharmacologic therapies mainly target a boost in dopamine production or inhibit its catabolism to alleviate the symptoms [28,44,94]. However, a cure is still definitely an unmet clinical need. The administration of antioxidants looks not to affect the course of PD, and the effect of statins (considered to act as anti-oxidants) is inconclusive, suggesting that redox stress might not be the primary responsible for DA reduction. Regarding these anti-inflammatory approaches (Table 1), non-steroidal anti-inflammatory drugs (NSAIDs) have been reported to lower up to 45% of PD disease risk [78]. Ibuprofen reduces up to a 27% PD incidence, possibly acting through the activation of PPAR γ [78] and decreasing DA turnover [76]. A similar result has been suggested for indomethacin: a weak agonist of PPAR [95]. Interventions working at the systemic level in decreasing systemic inflammation (see Section 4 for understanding the rationale) have also been considered. Moderate or vigorous physical activity is associated with a 34% reduced risk of PD, possibly by increasing BDNF release and DA synthesis [78] and/or decreasing the systemic inflammatory status [76,96]. Probiotics (lactobacilli, enterococci, bifidobacteria, yeasts, and various mixtures of beneficial bacteria) have been suggested to rebalance a PD-associated change in the microbiota composition, reducing the leaky gut and inflammation [96]. Tobacco (nicotine at a low dose, while at a high dose, it damages the BBB [76]) enhances TH expression and the DA level in PD through PPAR α . Still, the nicotine stimulation of the reward system explains Add [27,97]. Caffeine, theophylline, and theobromine increase DR activity in PD, decreasing the risk of developing the disease [76]. Based on the number of anti-inflammatory approaches and the results, even if partial, there is no doubt that PD is an inflammatory disease and that inflammation is a therapeutic target.

YPs have also been shown to participate in PD and inflammation. A comprehensive picture of PD, inflammation, DA, and YPs interplay could be recomposed by putting together the information belonging to different works. TSB is increased in the early stages of PD when it negatively correlates with the severity of symptoms and with less need for L-DOPA administration [98]. TSB has also been associated with the demise of DOPAn in models [99]. Moreover, in the brain parenchyma, the HMOX1 signal has been documented to be increased in specific areas (SNpc, regions of the neocortex with Lewy bodies) and cells (DOPAn, microglia, and astroglia) affected by the disease, and correlated with reduced inflammation, in agreement with the protective, anti-inflammatory action of bilirubin. On the contrary, (a) a TSB decrease even below the level of the controls has accompanied the progression of this disease to the most severe stages; (b) genetic polymorphisms of HMOX1 have also been shown to decrease BV and bilirubin production, correlating with the early onset of PD, stressing once again the importance of YPs and the modulation of inflammation to prevent or delay the pathology [100–105]. Based on this evidence, increased TSB, the increased presence of bilirubin degradation products in the urine, and an increased HMOX1 expression level have been suggested as potential markers for PD [35,103]. In models, bilirubin has been documented to induce TH enzyme and to increase DA levels in vivo (by NF κ B signaling pathway [106]). In turn, DA up-regulates the HMOX1 expression in vitro (endothelial cells [107]). Interestingly, DA may also inhibit the canonical, non-canonical, and α -syn-mediated activation of the NLRP3 inflammasome in vitro: a mechanism involved in DOPAn loss. Similar to the induction of HMOX1 and bilirubin increase in the early stages of PD, the inhibition of the inflammasome operated by DA might be interpreted as a tentative protective reaction to ongoing damage [108]. Notably, NRF2 is a primary regulator of cellular defense against oxidative stress and inflammation in DOPAn neuroprotection [18] and is one of the major signaling pathways of bilirubin protective action too [30] (Figure 2). On the other hand, the inhibitory effect of bilirubin on several steps of DA biology (tyrosine uptake; cAMP-stimulated DA synthesis, but not basal production; DA reuptake and cellular vesicular storage) has also been reported [109–112]. Part of these experiments have been conducted using toxic concentrations of bilirubin (70–140 μ M). This might explain the contradictory results. To this note, a harmful amount of bilirubin, as in severe neonatal hyperbilirubinemia, has been reported to trigger attention

deficit hyperactivity disorder (ADHD), schizophrenia (Sch), and autism spectrum disorder (ASD) in adult life, which are all known conditions in which DA is central, suggesting that a toxic amount of bilirubin could negatively affect one or more DA circuits (see later). It is worth mentioning that, in 2019, Chang et al. investigated the potential of idiopathic hyperbilirubinemia and Parkinson's disease to be associated with negative (lack of association) results [113]. In the Gunn rat, a spontaneous model of severe neonatal hyperbilirubinemia, DA evaluation in different parts of the brain gave contradictory results. DA concentration in the striatum, pons, medulla, and cerebellum was diminished based on Swenson [114] but was unaltered in the striatum and frontal cortex based on Shoko Miura [115]. This PD, DA, YPs, and inflammation interplay narrative needs experimental confirmation. Up until now, to the best of our knowledge, two of our works have supplied strong enough evidence on the protective role of supplementing tissue with bilirubin, counteracting inflammation. In organotypic brain cultures, models of PD (induced by rotenone) and inflammatory modulation (namely IL6, TNF α , and COX2), together with HMOX1 induction, precedes DA demise [81]; low (non-toxic) bilirubin supplementation fully protecting DOPAn specifically acts on TNF α [116]. Multiple pieces of evidence have supported these findings. NAC, as an anti-oxidant, is ineffective at protecting DOPAn. b) Infliximab (a clinic antibody neutralizing TNF α) administration to PD slices fully inhibits DOPAn demise. c) TNF α administration to healthy cultures induces a dose-dependent DOPAn loss identical to rotenone challenging. The question of whether this applies to other DA-related conditions has to be empirically evaluated. The following paragraph could serve as a background for understanding the potential of YPs and inflammation modulatory approaches in each condition. For completeness, it has to be remembered that the excessive activity of HMOX1 could lead to iron deposition and worsen the damage. DOPAn are not only intrinsically or highly sensitive to stress; however, DA itself might be easily oxidized, amplifying ongoing redox damage [1,42,103,117–126]. Thus, YPs and DA dysregulation might hypothetically result in a potent damaging couple in neurologic conditions. The data on YPs' potential in PD strongly suggests that the supplementation of exogenous bilirubin, for example, by nanobubbles, rather than inducing YPs's enzymes, could be a safer therapeutic approach.

3.2. Schizophrenia

Schizophrenia (Sch) is characterized by psychotic symptoms such as hallucinations, delusions, and disorganized speech, with negative symptoms including demotivation, reduced expressiveness, and cognitive deficiencies including impaired executive functioning, memory, and mental processing speed [127]. Positron emission tomography (PET) has shown an increase in DA synthesis and release in the dorsal striatum in individuals with Sch [128], supporting the role of DA in Sch as a long-prevailing hypothesis. This concept is further supported by a gene and transcriptome wide-association study of the postmortem brain, in which the gene encoding the dopamine DR2 was reported as a risk factor [129]. Psychotic symptoms tend to diminish in Sch patients who develop PD [130]. On the contrary, dopaminergic therapy induces psychosis in PD, and anti-psychotic therapy usually gives Parkinsonism as a side effect, highlighting the positive connection of DA with Sch (more DA, more Sch features) [131–133]. For this reason, the coexistence of both diseases challenges the treatment balance [134,135] (Figure 3).

The involvement of neuro-inflammation in Sch has been a topic of discussion. Epidemiological studies suggest that Sch (as well as ASD and ADHD) are developmental conditions on which neonatal stress can alter the physiologic synapse formation or neurotransmission development, with neurologic manifestations appearing in adult life. In line with this, pre-natal inflammation has been associated with an increased risk of Sch [136,137], with detrimental effects mediated by IL1 β , IL6, and TNF α [138–140]. Moreover, a pro-inflammatory status led by the gut was observed in patients with Sch, and the abundance of specific bacterial genera, *Succinivibrio* and *Corynebacterium*, has become significantly associated with the symptom severity of these diseases [141]. Concerning specifically TNF α ,

its serum level was found to be significantly higher before anti-psychotics treatment in first-episode drug-naïve Sch patients than in chronic patients and healthy controls, suggesting that an increase in $\text{TNF}\alpha$ was probably an early mechanism in the development of this disease [142]. This was supported by Lv et al., who reported a decrease in $\text{TNF}\alpha$ serum levels in chronic schizophrenia on long-term antipsychotics compared to healthy individuals [143] (Figure 2). Concerning the anti-inflammatory interventions in Sch, minocycline improved symptoms but not cognitive functions and aspirin has given contradictory results [144]. Statins (cholesterol-lowering agents, lowering also $\text{IL1}\beta$, IL6 , $\text{TNF}\alpha$, and CRP levels) may alleviate anxiety-like behavior in models of Sch by up-regulating the NMDA receptors, with improvements also reported in clinical trials, while in a study based on two subjects $\text{IFN}\gamma$ reduced the negative symptoms [144]. Omega-3 fatty acids (docosahexaenoic acid—DHA, eicosapentaenoic acid—EPA; anti-oxidant, suppressing $\text{NF}\kappa\beta$ and reducing the levels of IL1 , IL6 , and $\text{TNF}\alpha$ [145]) have provided contradictory results in Sch [144,146]. Possibly relevant to this review, celecoxib, as a selective blocker of COX2, looks to improve behavioral impairments in rodent models by reducing $\text{TNF}\alpha$ (Table 1). Its effects on DA levels have not been directly assessed, and its improvement is now referred to as the prevention of parvalbumin (PV) in calcium-binding GABAergic neuron loss [144]. Of relevance to this review, these neurons are one of the most hypothesized targets of bilirubin neurotoxicity in severe neonatal hyperbilirubinemia. This condition has also been suggested as a potential cause of certain diseases (see later).

A discrepancy exists in the findings on TSB levels in Sch. Some examples in the literature report a decrease in TSB [147], while others identify increased TSB levels in subjects without suspicion of genetic reasons for increased bilirubin in the blood [148]. A positive correlation (higher TSB, higher Sch incidence) also looked to be present in Gilbert's subjects, with a uridine glucuronidase (UGT1a1) defect that led to mild hyperbilirubinemia [149]. The genetic polymorphisms of the UGT1a1 enzyme in the Gilbert population could contribute even directly to DA concentration in the brain, with DA a substrate for the UGT1a mediated conjugation and dopamine glucuronide found in rat and mouse brain samples [150]. Even though it is still arguable, neonatal hyperbilirubinemia has been linked to schizophrenia in adult life, which is in agreement with the developmental hypotheses of Sch [151]. In agreement, the Gunn rat, the animal model of neonatal hyperbilirubinemia, has been used as a disease model as it presents Sch-like behavior [152,153]. In a recent publication, we reported that bilirubin mainly affected glutamate circuits in the brain, further supporting the potential link between higher bilirubin levels early after birth and the developmental hypothesis of schizophrenia [151]. The overexpression of HMOX1 in mice astrocytes resulted in DA augmentation and DR1 reduction and also has been linked with Sch-like features [154].

In summary, while increases in DA deficiency and $\text{TNF}\alpha$ cause motor PD, with the beneficial action of bilirubin, the excess DA with increased $\text{TNF}\alpha$ has recently been linked to Sch-related psychosis (Figures 2 and 3).

3.3. Attention Deficit Hyperactivity Disorder

With a prevalence of over 5%, attention deficit hyperactivity disorder (ADHD) is one of the most frequent disorders in child and adolescent psychiatry [155]. ADHD's clinical manifestations include a lack of attention, impulsivity, excessive motor activity, and hyperactivity [156]. Environmental etiological factors, including exposure to environmental toxins like organophosphate pesticides, polychlorinated biphenyls, and zinc, as well as prenatal and postnatal factors like maternal smoking and alcohol use, low birth weight, and premature birth, have all been linked to an increased risk for ADHD in past years [157]. The activation of TLR4 appears to be central [158]. A gene-by-environment interaction has been observed between the DAT1 (dopamine transporter 1) genotype and prenatal smoke exposure regarding hyperactivity—impulsivity [159,160]. DAT and dopamine receptor (DR1, DR2, DR4, and DR5) polymorphisms have been associated with an increase in ADHD [156,161–164]. In general, DA reduction and dysregulated DA neurotransmission,

particularly between the prefrontal and striate areas, are the molecular fingerprint of ADHD [165,166].

The potential involvement of neuroinflammation in ADHD has been discussed, and additional studies are needed for a more precise conclusion. Nevertheless, data on inflammation and ADHD are available. Neuroinflammation has been proposed as a risk factor for ADHD, as inflammation-related diseases such as atopic immune disorder, asthma, and rheumatoid arthritis, in addition to the perinatal conditions listed before, have been associated with the neuropsychiatric condition [167–171]. Particular emphasis on maternal inflammation in obesity and offspring neurodevelopmental disorders (ADHD and autism spectrum disorder) has been taken by Velda et al., who reported the activation of TLR, MAPK, NF κ B, and microglia, together with the epigenetic modulation of the brain environment which could have a critical role [158] (Figures 2 and 3). In agreement, a higher level of TNF-R1 (the receptor of TNF α) was found to be associated with ADHD [172], and a positive correlation between the TNF α serum level and hyperactivity-impulsivity was found in ADHD subjects [173]. Anand also explained ADHD as a pro-inflammatory condition (microgliosis, IL1 β , IL2, IL6, and TNF α), potentially affecting the development and/or functioning of the CNS, and finally reducing DA availability, in which perinatal infections are a relevant co-cause. The reduction in the neurotransmitter and the symptomatology could be mimicked in models by the injection of cytokines (IL1 β , IL2, and IL6) [168]. ADHD is also known as a neurodevelopmental condition. For example, the FOXP2 (forkhead box protein P2) gene, which controls DA and neurodevelopment in brain regions related to ADHD, was discovered to be strongly associated with ADHD in GWAS studies [174,175]. A reduction in the volume of the nucleus accumbens, amygdala, caudate, hippocampus, and putamen has been described in patients with ADHD [176].

Of relevance to this review, ADHD has been reported to be possibly induced by a post-natal exposition to the toxic bilirubin level. Chang-Ching Wei et al. reported a 2.48-fold increase in ADHD in the cohort of neonates presenting TSB levels requiring phototherapy and with longer admission days vs. the non-jaundiced cohort. The risk of neurologic sequelae was more remarkable for male, preterm, and low-birth-weight infants with neonatal jaundice [177]. A 30-year prospective study also unraveled that hyperbilirubinemia in early life could recreate a picture resembling the impairment of the frontostriatal network and the symptoms of the ADHD spectrum [178]. Indeed, bilirubin has been shown to have a dose-dependent inhibitory impact on the absorption of tyrosine: a precursor to dopamine [109]. Bilirubin has also been found to inhibit cAMP-stimulated dopamine synthesis and decrease its absorption and vesicular storage [110–112] (Figure 2).

Moreover, bilirubin-induced neurotoxicity is known to occur via the activation of microglia, astrogliosis, and the release of pro-inflammatory mediators, including TNF α , leading to developmental impairment, especially (but not limited to) basal ganglia and cerebellum [177,179], and as discussed in Sch, through the transcriptomic imprinting of genes that are involved mainly on glutamate neuro-transmission [151]. It suggests that at least high amounts of bilirubin may be dangerous in the context of ADHD. The effect of low (protective) doses of bilirubin (Gilbert-like of physiologic jaundice) or the modulation of YPs in the brain on ADHD is unknown. To complete the section, based on results in animal models, caffeine has also been suggested as an optional approach in this condition, acting on several mechanisms, including DA paired-adenosine receptors A_{2A}R, DAT level, and DA uptake at synapses. Of relevance for this review is among the reported targets of caffeine in ADHD, PKA, and PI3K, which are shared with bilirubin [180].

3.4. Autism Spectrum Disorder

During the past few decades, there have been indications of an increased prevalence of the neurodevelopmental disorder autism spectrum disorder (ASD) [181]. ASD has been linked to changes in dopamine signaling. Particularly changes in the mesocorticolimbic dopaminergic signaling system have been observed in autistic individuals, including decreased DA release in the prefrontal cortex and a decreased neuronal response in the

nucleus accumbens [182,183]. Several studies have reported that TNF α concentrations also increase in the blood, cerebrospinal fluid, or post-mortem brain samples of ASD subjects [184–187]. Moreover, children with autism display an elevated blood TNF α concentration correlating with symptom severity [188] (Figure 3). As previously mentioned, in agreement with the concept of the developmental disorder, an increased risk of ASD is present in infants exposed to maternal immune activation during pregnancy [158].

Also, for this condition, a negative correlation with TSB has been reported [179]. In a meta-analysis study (13 papers) on neonatal jaundice, the TSB level in the blood was associated with ASD in term infants but not in pre-terms, despite their moderate (10 mg/dL) TSB level [189]. Notably, similarly to ADHD, ASD is considered a neurodevelopmental disorder. The suggested adverse effect of bilirubin in DA, glutamate, and development biology could also be applied to ASD.

3.5. Depression

Depression (Dp, in figures) is one of the most serious mental disorders and is a major factor in the rising suicide rate in the 21st century, as a widespread mental disorder that jeopardizes the global population's physical and psychological health [190]. Recent research has revealed that the pathophysiology of depression might be caused by disturbances in DA regulatory networks, specifically the DOPAn populations implicated in motivation and reward. Some antidepressant drugs and brain stimulation therapies can impact the intricate DA system, with the selective inhibition of VTA DOPAn demonstrated to induce depression-like behavior in rodents [191].

According to clinical studies, peripheral inflammatory cytokines and their soluble receptors have been reported to be increased in the blood and cerebral fluid of individuals with major depression [192]. Cytokines may alter a range of dopamine neurotransmission processes, impairing vesicular DA packing and their release or increasing reuptake, which could interact to varying degrees to diminish basal ganglia dopamine neurotransmission [47]. Again, systemic inflammation (such as in metabolic syndrome) has been suggested to alter the brain DA circuit involved in depression and anhedonia, with the CNS modulation of IFN, IL1, IL6, TNF α , and COX2 levels, as a supposed critical role for TNF α and IL6 [193–195]. The results in models support this, where blocking two cytokines looks pivotal in reversing depressive symptoms [193]. Specifically, TNF α is known to damage the BBB, contributing to the development of the condition [196] (Figure 3). The pathophysiology of depression is thought to be mediated by TNF α signaling through the binding to its receptor TNFR1 [197]. In agreement, Infliximab, as a TNF α neutralizing antibody, can alleviate depressive symptoms in patients with treatment-resistant depression and elevated inflammatory markers [198]. Similarly, celecoxib, as a COX2 inhibitor, exhibited a satisfactory therapeutic effect in patients with this disease [199]. Aspirin improves anti-depressants' effect [195], possibly by MAPK modulation [200]. Minocycline application to depression has also resulted in inconclusive results [47,193], similar to statins and omega-3 fatty acids [193,195]. Combining ibuprofen and an anti-depressant gives disputed results [78,195] (Table 1).

Several studies have explored the association of bilirubin with depression. TSB has been reported to be significantly lower in major depressive disorder subjects and has been suggested as a result of bilirubin overconsumption to be an antioxidant [201]. Indeed, a low nocturnal bilirubin level is associated with winter seasonal depression [202].

Again, the picture of a decreased DA and increased inflammation, including TNF α , and reduced bilirubin, looks supportive of the possible beneficial anti-inflammatory action of pigment supply in depression.

3.6. Addiction

Addiction (Add) is regularly identified with the habitual nonmedical self-administration of drugs [203]. DA has typically been linked to the reinforcing effects of addictive substances and could play a significant role in initiating the neurobiological alterations connected to

addiction [165]. Different mechanisms of action on DA biology have been identified; however, in general, cocaine, amphetamines, opiates, alcohol, nicotine, and caffeine can elevate the extracellular neurotransmitter level in the Na core and Na shell in the ventral striatum region, which receives dopaminergic innervation from the ventral tegmental area (mesocorticolimbic DA system [204]), inducing a reinforcement effect [72,205], which is the basis of habit-forming [204]. To stress once more the link between systemic and CNS disease (see later), the same picture was documented in obese people with metabolic syndrome [206,207]. This supports the potential of shared neurologic mechanisms on the basis of addiction to drugs and “addiction” to palatable foods [208]. Specifically on the effects of abuse substances on DA biology, cocaine increases extracellular dopamine levels by preventing DAT from reabsorbing dopamine [203]. Amphetamines increase extracellular levels by releasing DA from vesicles [203]. Opiates activate the DA system by inhibiting GABA neurons which normally hold the dopamine neurons under inhibitory control [203]. Alcohol and cocaine increase DA release [76,209]. Similarly, nicotine induces Add by increasing the release of DA in Na while also promoting anxiety by inhibiting DA neurotransmission in Am [210].

It is believed that neuro-inflammation has a role in the neuronal changes that result from the long-term misuse of drug exposure [200], with morphological and functional alterations in the microglia and astrocytes characterizing the neuro-immune response to drugs in Add [211]. Microglia activation triggers cell migration to the site of injury, phagocytosis, and the production of pro-inflammatory mediators (IL1, IL6, and TNF α), as well as ROS and NOS that harm neurons [212–214]. Based on the translocator protein (TSPO—a marker of activated glia) neuro-imaging in methamphetamine users, the DA alterations are mediated by an increased inflammatory status (with IL6, TNF α release, and NOS and ROS generation), possibly through the increase in PPAR γ activity and protein levels, which results in increased release of IL1 β , IL6, and TNF α in the Na [212–215] (Figures 2 and 3). TNF α inhibits an increase in extracellular DA levels caused by methamphetamine while also activating plasmalemma and vesicular DA transporters, which could help prevent the drug dependence and neurotoxicity induced by methamphetamine [216]. Conversely, a pro-inflammatory CNS environment, with an increase in TNF α , could alleviate methamphetamine addiction [217]. Cannabis is largely considered therapeutic based on its anti-inflammatory effect [218], but increases in IL1 β , IL6, IL8, and TNF α have also been documented [219]. These discrepancies depend on the age at initiation of use [200]. For the potential benefits of other molecules, see Table 1 [25,27,97,220,221] (Table 1).

Very few data exist on YPs’ effects in Add. Interestingly, 6,7,4’-trihydroxyflavanone (THF), a flavone present in Leguminosae, might mitigate methamphetamine neurotoxicity by enhancing the NRF2 and PI3K signaling pathways, inducing HMOX1 expression and reducing oxidative stress and apoptosis in an in vitro neuronal model [222].

3.7. Ischemic Stroke

An ischemic stroke (IS) is characterized by a reduced brain blood and oxygen supply [223].

After an IS, a large DA release might be responsible for the neuronal damage brought on by ischemia through the activation of DR2 [224], followed by a smaller DA release for up to 3 days [225]. Subsequently, the genes that encode DR are down-regulated for at least one week [226]. One and two weeks after middle cerebral artery blockage, the striatum consistently exhibits decreased DR2 availability [227]. These findings can relate stroke to DA impairment, which could explain why people with a stroke have reduced implicit learning [228]. Moreover, the inhibition of dopamine DR1 and DR2 impaired motor skill recovery in an ischemic stroke model, implying the role of endogenous dopamine transmission in this rehabilitation activity (Figure 3).

Redox stress has been primarily implicated in DA-mediated neuronal sufferance. DR2 is a member of G protein-coupled receptors, which regulate cellular redox homeostasis [229]. Therefore, DR2 receptor agonists have neuroprotective potential by counteracting redox imbalance and mediating the anti-apoptotic process [230]. Indeed, the administration of DA DR2 agonists prevents neuron death by decreasing mitochondrial ROS

production [231]. Little information is available on inflammation in stroke and DA. In this respect, it has been reported that glutathione-mediated intrastratial DA production could reduce the cerebral infarction area and decrease inflammatory cytokines levels, including IL6 and TNF α [232], suggesting the damage-enhancing role for inflammation in the DA consequences of stroke. Parikh et al. recently reviewed the epidemiologic evidence of inflammation, autoimmunity, and infections on IS, reporting a possible correlation and partial result using anti-inflammatory approaches [233] (Figure 3 and Table 1).

The relationship between bilirubin and IS has been reported in some studies. Based on a Mendelian Randomization study, an inverse causal association between TSB and the risk of IS was shown [234]. Supporting these previous findings, a study on asymptomatic intracranial atherosclerosis as one of the IS risks found that participants with TSB lower than 12.30 $\mu\text{Mol/L}$ (0.72 mg/dL, with a normal range usually of 0.4 to 1.0 mg/dL) had an increased risk of developing cerebral atherosclerosis compared to those in the high (more than 12.3 $\mu\text{Mol/L}$ = 0.72 mg/dL) concentration group [235]. Of notice, only recently, low levels of TSB (in the lower normal range or below) have been proposed as a marker of risk for multiple neurologic conditions [35,80,236]. Of importance for this disease, the well-known anti-oxidant properties of bilirubin could be helpful in IS [237–239]. Nevertheless, considering the rise in DA at the site of the lesion, the hyperactivation of HMOX1 has to be avoided. As anticipated (see above), HMOX1 activation led to a rise in extracellular iron concentration, a potentially explosive mix. Again, the delivery of bilirubin but not the stimulation of YPs might be a reasonable approach.

3.8. Multiple Sclerosis

Multiple sclerosis (MS) is the most prevalent demyelinating condition affecting the central nervous system [240], in which DA is one of the most studied neurotransmitters in the modulation of neuro-immune interaction. Carandini et al. demonstrated that axonal damage in people with relapsing-remitting MS affects DA and norepinephrine neurotransmission [241]. Moreover, DA is also dysregulated in the peripheral immune system in MS [242]. In individuals with chronic progressive MS or relapsing-remitting MS, DA synthesis is decreased in activated lymphocytes [8]. In the peripheral blood mononuclear cells (T cells, B cells, and NK cells) of untreated relapsing-remitting MS patients, DR5 protein and mRNA levels are decreased [243].

Inflammatory cytokines seem to affect fatigue, as the neuropsychological symptoms of MS. Relapsing-remitting MS patients with fatigue were found to have greater plasma levels of IL6 and TNF α than the control group (relapsing-remitting MS patients without fatigue), and both cytokines concentrations were associated with fatigue severity [244]. A higher level of TNF α was also found in the CSF of MS patients compared to the controls [245]. IFN β is an immunomodulatory drug for relapsing-remitting MS, decreasing DA synthesis [246] (Figure 3).

In clinical settings, a decrease in TSB is frequently noted in MS patients [80]. As discussed in IS, a lower TSB has recently been hypothesized as a risk factor for neurologic diseases [30,35], and the potential biological explanations are described (see above). Studies on the animal model of MS, autoimmune encephalomyelitis (EAE), have described the protective function of bilirubin and YPs in this condition. Those studies demonstrated that the induction of the bilirubin-producing enzymes, HMOX1 and BLVR, reduced the symptoms of EAE, which acted on both the inflammatory and oxidative stress injury, protecting BBB integrity and reducing the invasion of inflammatory cells into the spinal cord [247–249]. Lately, more vital evidence of the anti-inflammatory effect of bilirubin was obtained by Kim et al., who injected bilirubin nanoparticles with the resulting mitigation of EAE progression by downregulating dendritic cells and Th17 production [250]. In conclusion, MS studies provide evidence that a low TSB may be a marker for the risk of disease progression, and the anti-inflammatory action of YPs might provide protection.

3.9. Late Neurologic Sequel of Neonatal Hyperbilirubinemia

Because in this review we have discussed the possible positive (restoring, alleviating, protective) interplay of YPs with inflammation and DA, it is mandatory to mention that severe neonatal hyperbilirubinemia has been supposed to correlate with DA neuropsychiatric syndromes, namely ADHD, Sch, and ASD. Despite this being a neglected area of research in the bilirubin field, some solid epidemiologic studies may be found [177,178,189,251]. For all these conditions, it has to be pointed out that, in severe neonatal hyperbilirubinemia, the beneficial effect of bilirubin turns upside down, with bilirubin becoming highly pro-inflammatory and interfering with brain development [153]. We have reported on the permanent transcriptomic alteration of genes involved with the aforementioned neurologic diseases [151], supporting the potential of a toxic amount of bilirubin in inducing neuropsychiatric conditions. Further studies are needed, especially if bilirubin should be considered as a therapeutic option.

3.10. Cerebral Palsy

Cerebral palsy (CP) is defined as a progressive injury to the developing CNS in children, leading to neurological and musculoskeletal abnormalities with severe motor disorders, including spasticity, dystonia, and immobility [252]. Prematurity, infections, and inflammation (TNF α / β , IL6, IL10) have been documented [253,254], and an increased TNF α level suggested a negative correlation with rehabilitation outcome [255].

The mesocortical DA circuit controls learned motor skills in humans [256] and models on which CREB/PKA signaling pathways have been suggested to be central [257]. The DA involvement in CP was supported by a study that assessed the association between successful rehabilitation (involving motor-learning abilities) and defined genetic polymorphisms on genes involved in DA biology (COMT—rs4680, DAT—rs28363170, DRD1—rs4532, DRD2—rs1800497, DRD3—rs6280) on 33 patients with spastic unilateral CP [258]. The association between DA genetic variants and motor skills in CP infants was corroborated in 498 extremely low-birthweight infants [259]. Indeed, DA is a treatment in children with dystonia and CP [252].

CP is also a known feature of severe neonatal hyperbilirubinemia, with bilirubin reported to damage the basal ganglia and cerebellum [260,261]. Notably, the cerebellum is also a central region in controlling body movements. Currently, there are no studies that assess the effect of severe neonatal hyperbilirubinemia and DA biology in CP. Nevertheless, as discussed in Section 3.1, the DA level was found to decrease in the striatum, pons, medulla, and cerebellum of the Gunn rat [114]. Together with the known pro-inflammatory effect of sustained TSB, and the shared target signaling pathways, this suggests the involvement of high (toxic) TSB, inflammation, and DA dysregulation in the development of neonatal CP.

3.11. Other Conditions Linked to DA Biology

For the other conditions due to alterations in the DA system, such as the restless leg syndrome, minimal hepatic encephalopathy, pain potentially inducing anhedonia-like behavior, and the emotional and cognitive deficits involving the mesocortical dopaminergic system, no data support a discussion on DA, inflammation or the interplay of YPs.

Table 1. Interventions with beneficial effects on DA-related diseases.

Types	Drugs/Compounds	DA-Related Disease	Ref.
Anti-inflammatory drugs	NSAIDs	PD	[78]
	Ibuprofen	PD, Dp	[78,195]
	Indomethacin	PD	[95]
	Celecoxib	Sch	[144]
	Aspirin	Dp, Add	[95]

Table 1. Cont.

Types	Drugs/Compounds	DA-Related Disease	Ref.
	Minocycline	Sch	[144]
	Statins	Sch	[144]
	IFN β	MS	[246]
	IFN γ	Sch	[144]
	TNF α antagonists	Dp	[198]
Physical activity	-	PD	[76,78]
Nutritional interventions	Omega-3 fatty acids	Sch, Dp	[174,193,195]
	Oleoylethanolamide (OEA) and palmitoylethanolamide (PEA)	Add	[25,27,220]
	Probiotics	PD	[96]
Others	Tobacco	PD	[76]
	Caffeine	PD, ADHD	[76,180]
	Theophylline	PD	[76]
	Theobromine	PD	[76]
YPs modulation		PD, Dp, MS	[44,80,116]

NSAIDs: non-steroidal anti-inflammatory drugs; IFN: interferon; TNF: tumor necrosis factor; YPs: yellow players; PD: Parkinson's disease; Dp: depression; Sch: schizophrenia; Add: addiction; MS: multiple sclerosis; ADHD: Attention Deficit Hyperactivity Disorder.

4. DA, Systemic Inflammation and the Brain

A recent vision on neurologic conditions has suggested that the brain is not as isolated as it has been said and that systemic stimuli may either affect or even trigger neurologic diseases.

DRs are expressed by T-cells, dendritic cells, B cells, osteoclasts, neutrophils, eosinophils, monocyte, and macrophages [11,22], which link together DA and systemic inflammation (infections, sepsis, colitis, rheumatoid arthritis, tumors, liver injury, metabolic syndrome, inflammatory bowel diseases, Chron disease, diabetes, etc.) to neurologic conditions (Dp, PD, ASD, etc.) [11,14]. Moreover, DA is also synthesized by the mesenteric organs (approximately 45% of all the DA body production), the sympathetic nerves, the gut microbiota (with *Prevotella*, *Bacteroides*, *Lactobacillus*, *Bifidobacterium*, *Clostridium*, *Enterococcus*, and *Ruminococcus* that may modulate DA availability, DA then transported to the brain thanks to the vagus nerve [262,263]), the adrenal medulla, and the kidney [10,264]. Notably, Klingelhoefer and Reichmann even suggested that the pre-motor symptoms (olfactory impairment and constipation) could be explained by the origin of the disease in the enteric nervous system (ENS) before propagating via the parasympathetic trans-synaptic cell-to-cell of alpha-synuclein through the nervus vagus, playing a Trojan horse role [21]. Because the gut is a route for the environment to the body, this may strongly support the potential of environmental substances as triggers of neurologic diseases.

The connection between systemic and brain status in diseases is also supported by metabolic syndrome data. Metabolic syndrome is a growing world problem, especially among young people. In the metabolic syndrome, a diet rich in calories and fat and poor in fibers; microbiota dysbiosis; and low physical activity induce a low pro-inflammatory systemic status, increasing the risk of developing diabetes, cardiovascular diseases (CVD), and CVD-related problems (stroke, ischemia, etc.). Similarly, dietary constituents and a sedentary lifestyle have repeatedly been shown to influence the plasma levels of DA [264,265], strengthening the link between lifestyle, DA, inflammation, the gut, and the brain. Special emphasis on obesity and maternal inflammation concerning offspring neurodevelopmental disorders has been taken by Velda et al. [158]. A link between adiposity and depression

has also been suggested and motivated again through an increased systemic inflammatory situation and the increased risk of coronary heart disease [266,267].

While the potential interaction between DA and YPs in the CNS has been discussed above, it is worth mentioning that bilirubin is known to act at the systemic level [30,31,35]. Epidemiologic data (on more than 100,000 subjects belonging to different areas of the world) support bilirubin as an important physiologic modulator of chronic inflammation in metabolic syndrome and diabetes. Gilbert subjects are usually lean and resistant to metabolic syndrome and diabetes type 2. BLVR induction and bilirubin increase have been reported to reduce hepatic steatosis [268,269], possibly through bilirubin binding to PPAR, as a central signaling pathway in obesity, which induces the fat-burning genes *Cpt1*, *Ucp1*, and *Adrb3* (carnitine palmitoyltransferase 1A, uncoupling protein 1, adrenoceptor beta 3) [35,270]. Also reported is reduced inflammation in the adipose samples, pancreas, and liver, with decreased $\text{TNF}\alpha$ and $\text{IL1}\beta$ [271]. The induction of HMOX1 with even a marginal elevation in bilirubin is protective against diabetes type 2 and vascular complications, including retinopathy, nephropathy, and CVD [272], increasing insulin sensitivity, IRS2 expression and regulating its signaling [268,269,273].

5. Future Perspective

The inflammation, dysfunctions/diseases of DA biology, and YPs share multiple molecular mechanisms and signaling pathways, both systemically and at the CNS level. Due to its recognized potential for health, the modulation of YPs is a hot research topic.

YPs, and especially HMOX1, may be modulated, giving the possibility to increase bilirubin levels in the blood [36,274–276] and offering a potential clinic application for DA-related pathologies.

The BBB is a limitation to the pharmacologic modulation of YPs in the brain; however, new molecules with better CNS bioavailability can be developed. However, HMOX1 hyperactivation in the brain might be deleterious due to iron deposition with the worsening of the disease.

Bilirubin is known to cross the BBB by passive diffusion from the systemic circulation [277]; however, controlling the level of this pigment that reaches the parenchyma after its systemic increase is not easy, if not impossible. Nevertheless, the documented anti-inflammatory benefits of bilirubin have attracted interest in nanomedicine, and bilirubin-loaded nano-delivery strategies are under investigation with promising results in other contexts (tumors, metabolic syndrome, etc.) [37,278–283]. A multidisciplinary (material science, physics, biology, medicine, etc.) approach is essential.

6. Conclusions

While in certain neurologic and neuropsychiatric conditions, the connections between DA, YPs, and inflammation are still scarce and need additional studies, and in Sch, bilirubin appears dangerous irrespective of its level, in the larger part of the CNS diseases here reported as well as on their systemic inflammatory component, bilirubin seems to be an endogenous candidate biomolecule for improving or preventing DA related neurologic dysfunctions, supporting the potential for its therapeutic usage.

Author Contributions: Conceptualization, S.J., C.D.V., C.T. and S.G.; data curation, S.J., C.D.V., C.T. and S.G.; writing—original draft preparation, S.J., C.D.V., C.T. and S.G.; writing—review and editing, C.T. and S.G. All authors have read and agreed to the published version of the manuscript.

Funding: This work was partly funded by an in-house grant from the Italian Liver Foundation (C.D.V, S.G., C.T.) and partly by a post-doctoral fellowship supported by the National Research and Innovation Agency of Indonesia (S.J.).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: All data are contained within the article.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Chinta, S.J.; Andersen, J.K. Dopaminergic neurons. *Int. J. Biochem. Cell Biol.* **2005**, *37*, 942–946. [\[CrossRef\]](#) [\[PubMed\]](#)
- Macedo-Lima, M.; Remage-Healey, L. Dopamine Modulation of Motor and Sensory Cortical Plasticity among Vertebrates. *Integr. Comp. Biol.* **2021**, *61*, 316–336. [\[CrossRef\]](#) [\[PubMed\]](#)
- Schultz, W. Multiple dopamine functions at different time courses. *Annu. Rev. Neurosci.* **2007**, *30*, 259–288. [\[CrossRef\]](#)
- Furuyashiki, T. Roles of dopamine and inflammation-related molecules in behavioral alterations caused by repeated stress. *J. Pharmacol. Sci.* **2012**, *120*, 63–69. [\[CrossRef\]](#) [\[PubMed\]](#)
- Nieoullon, A. Dopamine and the regulation of cognition and attention. *Prog. Neurobiol.* **2002**, *67*, 53–83. [\[CrossRef\]](#)
- Carmichael, K.; Sullivan, B.; Lopez, E.; Sun, L.; Cai, H. Diverse midbrain dopaminergic neuron subtypes and implications for complex clinical symptoms of Parkinson’s disease. *Ageing Neurodegener. Dis.* **2021**, *1*. [\[CrossRef\]](#)
- Neumann, J.; Hofmann, B.; Dhein, S.; Gergs, U. Role of Dopamine in the Heart in Health and Disease. *Int. J. Mol. Sci.* **2023**, *24*, 5042. [\[CrossRef\]](#)
- Cosentino, M.; Zaffaroni, M.; Marino, F.; Bombelli, R.; Ferrari, M.; Rasini, E.; Lecchini, S.; Ghezzi, A.; Frigo, G. Catecholamine production and tyrosine hydroxylase expression in peripheral blood mononuclear cells from multiple sclerosis patients: Effect of cell stimulation and possible relevance for activation-induced apoptosis. *J. Neuroimmunol.* **2002**, *133*, 233–240. [\[CrossRef\]](#)
- Channer, B.; Matt, S.M.; Nickoloff-Bybel, E.A.; Pappa, V.; Agarwal, Y.; Wickman, J.; Gaskill, P.J. Dopamine, Immunity, and Disease. *Pharmacol. Rev.* **2023**, *75*, 62–158. [\[CrossRef\]](#)
- Leite, F.; Ribeiro, L. Dopaminergic Pathways in Obesity-Associated Inflammation. *J. Neuroimmune Pharmacol.* **2020**, *15*, 93–113. [\[CrossRef\]](#)
- Li, M.; Zhou, L.; Sun, X.; Yang, Y.; Zhang, C.; Wang, T.; Fu, F. Dopamine, a co-regulatory component, bridges the central nervous system and the immune system. *Biomed. Pharmacother.* **2022**, *145*, 112458. [\[CrossRef\]](#)
- Albanese, A.; Altavista, M.C.; Rossi, P. Organization of central nervous system dopaminergic pathways. *J. Neural Transm. Suppl.* **1986**, *22*, 3–17. [\[PubMed\]](#)
- Rangel-Barajas, C.; Coronel, I.; Florán, B. Dopamine Receptors and Neurodegeneration. *Aging Dis.* **2015**, *6*, 349–368. [\[CrossRef\]](#) [\[PubMed\]](#)
- Carandini, T.; Cercignani, M.; Galimberti, D.; Scarpini, E.; Bozzali, M. The distinct roles of monoamines in multiple sclerosis: A bridge between the immune and nervous systems? *Brain Behav. Immun.* **2021**, *94*, 381–391. [\[CrossRef\]](#)
- Martin-de-Pablos, A.; Córdoba-Fernández, A.; Fernández-Espejo, E. Analysis of neurotrophic and antioxidant factors related to midbrain dopamine neuronal loss and brain inflammation in the cerebrospinal fluid of the elderly. *Exp. Gerontol.* **2018**, *110*, 54–60. [\[CrossRef\]](#)
- Scherman, D.; Desnos, C.; Darchen, F.; Pollak, P.; Javoy-Agid, F.; Agid, Y. Striatal dopamine deficiency in parkinson’s disease: Role of aging. *Ann. Neurol.* **1989**, *26*, 551–557. [\[CrossRef\]](#)
- Fearnley, J.M.; Lees, A.J. Ageing and Parkinson’s disease: Substantia nigra regional selectivity. *Brain* **1991**, *114*, 2283–2301. [\[CrossRef\]](#)
- Marchetti, B.; Giachino, C.; Tirolo, C.; Serapide, M.F. “Reframing” dopamine signaling at the intersection of glial networks in the aged Parkinsonian brain as innate Nrf2/Wnt driver: Therapeutical implications. *Aging Cell* **2022**, *21*, e13575. [\[CrossRef\]](#)
- Furuyashiki, T.; Kitaoka, S. Neural mechanisms underlying adaptive and maladaptive consequences of stress: Roles of dopaminergic and inflammatory responses. *Psychiatry Clin. Neurosci.* **2019**, *73*, 669–675. [\[CrossRef\]](#) [\[PubMed\]](#)
- Vidal, P.M.; Pacheco, R. Targeting the Dopaminergic System in Autoimmunity. *J. Neuroimmune Pharmacol. Off. J. Soc. NeuroImmune Pharmacol.* **2020**, *15*, 57–73. [\[CrossRef\]](#)
- Klingelhoefer, L.; Reichmann, H. Pathogenesis of Parkinson disease—the gut-brain axis and environmental factors. *Nat. Rev. Neurol.* **2015**, *11*, 625–636. [\[CrossRef\]](#) [\[PubMed\]](#)
- Feng, Y.; Lu, Y. Immunomodulatory Effects of Dopamine in Inflammatory Diseases. *Front. Immunol.* **2021**, *12*, 663102. [\[CrossRef\]](#)
- Rauf, A.; Badoni, H.; Abu-Izneid, T.; Olatunde, A.; Rahman, M.M.; Painuli, S.; Semwal, P.; Wilairatana, P.; Mubarak, M.S. Neuroinflammatory Markers: Key Indicators in the Pathology of Neurodegenerative Diseases. *Molecules* **2022**, *27*, 3194. [\[CrossRef\]](#) [\[PubMed\]](#)
- Felger, J.C.; Treadway, M.T. Inflammation Effects on Motivation and Motor Activity: Role of Dopamine. *Neuropsychopharmacology* **2017**, *42*, 216–241. [\[CrossRef\]](#)
- Nolan, Y.M.; Sullivan, A.M.; Toulouse, A. Parkinson’s disease in the nuclear age of neuroinflammation. *Trends Mol. Med.* **2013**, *19*, 187–196. [\[CrossRef\]](#)
- Xia, Q.-P.; Cheng, Z.-Y.; He, L. The modulatory role of dopamine receptors in brain neuroinflammation. *Int. Immunopharmacol.* **2019**, *76*, 105908. [\[CrossRef\]](#)
- Melis, M.; Carta, G.; Pistis, M.; Banni, S. Physiological Role of Peroxisome Proliferator-Activated Receptors Type Alpha on Dopamine Systems. *CNS Neurol. Disord.-Drug Targets-CNS Neurol. Disord.* **2013**, *12*, 70–77. [\[CrossRef\]](#)
- Pajares, M.; Rojo, A.I.; Manda, G.; Boscá, L.; Cuadrado, A. Inflammation in Parkinson’s Disease: Mechanisms and Therapeutic Implications. *Cells* **2020**, *9*, 1687. [\[CrossRef\]](#)

29. Vitek, L.; Ostrow, J.D. Bilirubin Chemistry and Metabolism; Harmful and Protective Aspects. *Curr. Pharm. Des.* **2009**, *15*, 2869–2883. [CrossRef] [PubMed]
30. Wagner, K.-H.; Wallner, M.; Mölzer, C.; Gazzin, S.; Bulmer, A.C.; Tiribelli, C.; Vitek, L. Looking to the horizon: The role of bilirubin in the development and prevention of age-related chronic diseases. *Clin. Sci. Lond. Engl.* 1979 **2015**, *129*, 1–25. [CrossRef]
31. Gazzin, S.; Vitek, L.; Watchko, J.; Shapiro, S.M.; Tiribelli, C. A Novel Perspective on the Biology of Bilirubin in Health and Disease. *Trends Mol. Med.* **2016**, *22*, 758–768. [CrossRef]
32. Chen, J.; Tu, Y.; Moon, C.; Nagata, E.; Ronnett, G.V. Heme oxygenase-1 and heme oxygenase-2 have distinct roles in the proliferation and survival of olfactory receptor neurons mediated by cGMP and bilirubin, respectively. *J. Neurochem.* **2003**, *85*, 1247–1261. [CrossRef]
33. Park, J.-S.; Nam, E.; Lee, H.-K.; Lim, M.H.; Rhee, H.-W. In Cellulo Mapping of Subcellular Localized Bilirubin. *ACS Chem. Biol.* **2016**, *11*, 2177–2185. [CrossRef] [PubMed]
34. Takeda, T.; Mu, A.; Tai, T.T.; Kitajima, S.; Taketani, S. Continuous de novo biosynthesis of haem and its rapid turnover to bilirubin are necessary for cytoprotection against cell damage. *Sci. Rep.* **2015**, *5*, 10488. [CrossRef] [PubMed]
35. Creeden, J.F.; Gordon, D.M.; Stec, D.E.; Hinds, T.D. Bilirubin as a Metabolic Hormone: The Physiological Relevance of Low Levels. *Am. J. Physiol.-Endocrinol. Metab.* **2020**, *320*, E191–E207. [CrossRef] [PubMed]
36. Vitek, L.; Bellarosa, C.; Tiribelli, C. Induction of Mild Hyperbilirubinemia: Hype or Real Therapeutic Opportunity? *Clin. Pharmacol. Ther.* **2019**, *106*, 568–575. [CrossRef]
37. Hinds, T.D.J.; Creeden, J.F.; Gordon, D.M.; Stec, D.F.; Donald, M.C.; Stec, D.E. Bilirubin Nanoparticles Reduce Diet-Induced Hepatic Steatosis, Improve Fat Utilization, and Increase Plasma β -Hydroxybutyrate. *Front. Pharmacol.* **2020**, *11*, 594574. [CrossRef]
38. Schipper, H.M.; Song, W.; Tavitian, A.; Cressatti, M. The sinister face of heme oxygenase-1 in brain aging and disease. *Prog. Neurobiol.* **2019**, *172*, 40–70. [CrossRef]
39. Abraham, N.G.; Kappas, A. Pharmacological and clinical aspects of heme oxygenase. *Pharmacol. Rev.* **2008**, *60*, 79–127. [CrossRef]
40. Gozzelino, R. The Pathophysiology of Heme in the Brain. Available online: <https://www.eurekaselect.com/135089/article> (accessed on 27 July 2020).
41. Maines, M.D. New Insights into Biliverdin Reductase Functions: Linking Heme Metabolism to Cell Signaling. *Physiology* **2005**, *20*, 382–389. [CrossRef]
42. Nitti, M.; Piras, S.; Brondolo, L.; Marinari, U.M.; Pronzato, M.A.; Furfaro, A.L. Heme Oxygenase 1 in the Nervous System: Does It Favor Neuronal Cell Survival or Induce Neurodegeneration? *Int. J. Mol. Sci.* **2018**, *19*, 2260. [CrossRef] [PubMed]
43. Ryter, S.W.; Alam, J.; Choi, A.M.K. Heme oxygenase-1/carbon monoxide: From basic science to therapeutic applications. *Physiol. Rev.* **2006**, *86*, 583–650. [CrossRef] [PubMed]
44. Jayanti, S.; Moretti, R.; Tiribelli, C.; Gazzin, S. Bilirubin: A Promising Therapy for Parkinson’s Disease. *Int. J. Mol. Sci.* **2021**, *22*, 6223. [CrossRef]
45. Fuxe, K. Evidence for the existence of monoamine neurons in the central nervous system. *Z. Für Zellforsch. Mikrosk. Anat.* **1965**, *65*, 573–596. [CrossRef]
46. Wise, R.A. Dopamine, learning and motivation. *Nat. Rev. Neurosci.* **2004**, *5*, 483–494. [CrossRef]
47. Felger, J.C. The Role of Dopamine in Inflammation-Associated Depression: Mechanisms and Therapeutic Implications. *Curr. Top. Behav. Neurosci.* **2017**, *31*, 199–219. [CrossRef]
48. Alcaro, A.; Huber, R.; Panksepp, J. Behavioral functions of the mesolimbic dopaminergic system: An affective neuroethological perspective. *Brain Res. Rev.* **2007**, *56*, 283–321. [CrossRef]
49. Ayano, G. Dopamine: Receptors, Functions, Synthesis, Pathways, Locations and Mental Disorders: Review of Literatures. *J. Ment. Disord. Treat.* **2016**, *2*, 2. [CrossRef]
50. Reid, J.L. Dopaminergic Pathways and their Pathophysiological Significance. *Clin. Sci. Mol. Med.* **1977**, *53*, 303–306. [CrossRef] [PubMed]
51. Hisahara, S.; Shimohama, S. Dopamine receptors and Parkinson’s disease. *Int. J. Med. Chem.* **2011**, *2011*, 403039. [CrossRef]
52. Grahm, J.A.; Parkinson, J.A.; Owen, A.M. The role of the basal ganglia in learning and memory: Neuropsychological studies. *Behav. Brain Res.* **2009**, *199*, 53–60. [CrossRef] [PubMed]
53. Liang, L.; DeLong, M.R.; Papa, S.M. Inversion of Dopamine Responses in Striatal Medium Spiny Neurons and Involuntary Movements. *J. Neurosci.* **2008**, *28*, 7537–7547. [CrossRef] [PubMed]
54. Griffiths, P.D.; Perry, R.H.; Crossman, A.R. A detailed anatomical analysis of neurotransmitter receptors in the putamen and caudate in Parkinson’s disease and Alzheimer’s disease. *Neurosci. Lett.* **1994**, *169*, 68–72. [CrossRef] [PubMed]
55. Schrimsher, G.W.; Billingsley, R.L.; Jackson, E.F.; Moore, B.D. Caudate Nucleus Volume Asymmetry Predicts Attention-Deficit Hyperactivity Disorder (ADHD) Symptomatology in Children. *J. Child Neurol.* **2002**, *17*, 877–884. [CrossRef]
56. Fusar-Poli, P.; Meyer-Lindenberg, A. Striatal Presynaptic Dopamine in Schizophrenia, Part II: Meta-Analysis of [^{18}F / ^{11}C]-DOPA PET Studies. *Schizophr. Bull.* **2013**, *39*, 33–42. [CrossRef]
57. Walker, F.O. Huntington’s disease. *Lancet Lond. Engl.* **2007**, *369*, 218–228. [CrossRef]
58. Douglas, P.K.; Gutman, B.; Anderson, A.; Larios, C.; Lawrence, K.E.; Narr, K.; Sengupta, B.; Cooray, G.; Douglas, D.B.; Thompson, P.M.; et al. Hemispheric brain asymmetry differences in youths with attention-deficit/hyperactivity disorder. *NeuroImage Clin.* **2018**, *18*, 744–752. [CrossRef]

59. Weinstein, J.J.; Chohan, M.O.; Slifstein, M.; Kegeles, L.S.; Moore, H.; Abi-Dargham, A. Pathway-Specific Dopamine Abnormalities in Schizophrenia. *Biol. Psychiatry* **2017**, *81*, 31–42. [\[CrossRef\]](#)
60. Maller, J.J.; Thaveenthiran, P.; Thomson, R.H.; McQueen, S.; Fitzgerald, P.B. Volumetric, cortical thickness and white matter integrity alterations in bipolar disorder type I and II. *J. Affect. Disord.* **2014**, *169*, 118–127. [\[CrossRef\]](#)
61. Telzer, E.H. Dopaminergic reward sensitivity can promote adolescent health: A new perspective on the mechanism of ventral striatum activation. *Dev. Cogn. Neurosci.* **2016**, *17*, 57–67. [\[CrossRef\]](#)
62. Baik, J.-H. Stress and the dopaminergic reward system. *Exp. Mol. Med.* **2020**, *52*, 1879–1890. [\[CrossRef\]](#)
63. Serafini, R.A.; Pryce, K.D.; Zachariou, V. The Mesolimbic Dopamine System in Chronic Pain and Associated Affective Comorbidities. *Biol. Psychiatry* **2020**, *87*, 64–73. [\[CrossRef\]](#) [\[PubMed\]](#)
64. Yang, S.; Boudier-Rev  ret, M.; Choo, Y.J.; Chang, M.C. Association between Chronic Pain and Alterations in the Mesolimbic Dopaminergic System. *Brain Sci.* **2020**, *10*, 701. [\[CrossRef\]](#) [\[PubMed\]](#)
65. Floresco, S.B.; Magyar, O. Mesocortical dopamine modulation of executive functions: Beyond working memory. *Psychopharmacology* **2006**, *188*, 567–585. [\[CrossRef\]](#) [\[PubMed\]](#)
66. Berridge, K.C.; Kringelbach, M.L. Pleasure systems in the brain. *Neuron* **2015**, *86*, 646–664. [\[CrossRef\]](#) [\[PubMed\]](#)
67. Weinberger, D.R.; Berman, K.F.; Chase, T.N. Mesocortical dopaminergic function and human cognition. *Ann. N. Y. Acad. Sci.* **1988**, *537*, 330–338. [\[CrossRef\]](#)
68. Hadland, K.A.; Rushworth, M.F.S.; Gaffan, D.; Passingham, R.E. The effect of cingulate lesions on social behaviour and emotion. *Neuropsychologia* **2003**, *41*, 919–931. [\[CrossRef\]](#)
69. Habibi, M. Dopamine Receptors. In *Encyclopedia of Movement Disorders*; Kompoliti, K., Metman, L.V., Eds.; Academic Press: Oxford, UK, 2010; pp. 326–329. ISBN 978-0-12-374105-9.
70. Katolikova, N.; Gainetdinov, R.R. Dopamine System. In *Encyclopedia of Molecular Pharmacology*; Offermanns, S., Rosenthal, W., Eds.; Springer International Publishing: Cham, Switzerland, 2021; pp. 554–560. ISBN 978-3-030-57401-7.
71. Lyons, D.J.; Hellysaz, A.; Broberger, C. Prolactin Regulates Tuberoinfundibular Dopamine Neuron Discharge Pattern: Novel Feedback Control Mechanisms in the Lactotrophic Axis. *J. Neurosci.* **2012**, *32*, 8074–8083. [\[CrossRef\]](#)
72. Robbins, T.W. Dopamine and cognition. *Curr. Opin. Neurol.* **2003**, *16* (Suppl. 2), S1–S2. [\[CrossRef\]](#)
73. Dorsey, E.R.; Bloem, B.R. The Parkinson Pandemic-A Call to Action. *JAMA Neurol.* **2018**, *75*, 9–10. [\[CrossRef\]](#)
74. Surmeier, D.J.; Obeso, J.A.; Halliday, G.M. Selective neuronal vulnerability in Parkinson disease. *Nat. Rev. Neurosci.* **2017**, *18*, 101–113. [\[CrossRef\]](#) [\[PubMed\]](#)
75. Hirano, S. Clinical implications for dopaminergic and functional neuroimage research in cognitive symptoms of Parkinson’s disease. *Mol. Med.* **2021**, *27*, 40. [\[CrossRef\]](#) [\[PubMed\]](#)
76. Adams, J.D. Possible causes of Parkinson’s disease. *Front. Biosci. Landmark Ed.* **2021**, *26*, 387–394. [\[CrossRef\]](#)
77. Chen, H.; Jacobs, E.; Schwarzschild, M.A.; McCullough, M.L.; Calle, E.E.; Thun, M.J.; Ascherio, A. Nonsteroidal antiinflammatory drug use and the risk for Parkinson’s disease. *Ann. Neurol.* **2005**, *58*, 963–967. [\[CrossRef\]](#) [\[PubMed\]](#)
78. Ascherio, A.; Schwarzschild, M.A. The epidemiology of Parkinson’s disease: Risk factors and prevention. *Lancet Neurol.* **2016**, *15*, 1257–1272. [\[CrossRef\]](#) [\[PubMed\]](#)
79. Kannarkat, G.T.; Boss, J.M.; Tansey, M.G. The Role of Innate and Adaptive Immunity in Parkinson’s Disease. *J. Park. Dis.* **2013**, *3*, 493–514. [\[CrossRef\]](#)
80. Jayanti, S.; Moretti, R.; Tiribelli, C.; Gazzin, S. Bilirubin and inflammation in neurodegenerative and other neurological diseases. *Neuroinflammation* **2020**, *7*, 92–108. [\[CrossRef\]](#)
81. Dal Ben, M.; Bongiovanni, R.; Tuniz, S.; Fioriti, E.; Tiribelli, C.; Moretti, R.; Gazzin, S. Earliest Mechanisms of Dopaminergic Neurons Sufferance in a Novel Slow Progressing Ex Vivo Model of Parkinson Disease in Rat Organotypic Cultures of Substantia Nigra. *Int. J. Mol. Sci.* **2019**, *20*, 2224. [\[CrossRef\]](#)
82. Tansey, M.G.; McCoy, M.K.; Frank-Cannon, T.C. Neuroinflammatory mechanisms in Parkinson’s disease: Potential environmental triggers, pathways, and targets for early therapeutic intervention. *Exp. Neurol.* **2007**, *208*, 1–25. [\[CrossRef\]](#)
83. Menza, M.; Dobkin, R.D.; Marin, H.; Mark, M.H.; Gara, M.; Bienfait, K.; Dicke, A.; Kusnekov, A. The role of inflammatory cytokines in cognition and other non-motor symptoms of Parkinson’s disease. *Psychosomatics* **2010**, *51*, 474–479. [\[CrossRef\]](#)
84. Rocha, N.P.; Teixeira, A.L.; Scalzo, P.L.; Barbosa, I.G.; de Sousa, M.S.; Morato, I.B.; Vieira, E.L.M.; Christo, P.P.; Palot  s, A.; Reis, H.J. Plasma levels of soluble tumor necrosis factor receptors are associated with cognitive performance in Parkinson’s disease. *Mov. Disord. Off. J. Mov. Disord. Soc.* **2014**, *29*, 527–531. [\[CrossRef\]](#) [\[PubMed\]](#)
85. Imamura, K.; Hishikawa, N.; Sawada, M.; Nagatsu, T.; Yoshida, M.; Hashizume, Y. Distribution of major histocompatibility complex class II-positive microglia and cytokine profile of Parkinson’s disease brains. *Acta Neuropathol.* **2003**, *106*, 518–526. [\[CrossRef\]](#) [\[PubMed\]](#)
86. Jenner, P. Oxidative stress in Parkinson’s disease. *Ann. Neurol.* **2003**, *53* (Suppl. 3), S26–S38. [\[CrossRef\]](#)
87. Bosco, D.A.; Fowler, D.M.; Zhang, Q.; Nieva, J.; Powers, E.T.; Wentworth, P.; Lerner, R.A.; Kelly, J.W. Elevated levels of oxidized cholesterol metabolites in Lewy body disease brains accelerate α -synuclein fibrilization. *Nat. Chem. Biol.* **2006**, *2*, 249–253. [\[CrossRef\]](#)
88. Nishimura, M.; Mizuta, I.; Mizuta, E.; Yamasaki, S.; Ohta, M.; Kaji, R.; Kuno, S. Tumor necrosis factor gene polymorphisms in patients with sporadic Parkinson’s disease. *Neurosci. Lett.* **2001**, *311*, 1–4. [\[CrossRef\]](#)

89. Sun, X.; He, Y.; Shen, Z.; Chen, L. TNF- α (G308A) Polymorphism and Vascular Parkinson Complicated with Pulmonary Infection. *J. Coll. Physicians Surg.—Pak. JCPSP* **2022**, *32*, 860–863. [\[CrossRef\]](#)
90. Lutters, B.; Foley, P.; Koehler, P.J. The centennial lesson of encephalitis lethargica. *Neurology* **2018**, *90*, 563–567. [\[CrossRef\]](#) [\[PubMed\]](#)
91. Limphaibool, N.; Iwanowski, P.; Holstad, M.J.V.; Kobylarek, D.; Kozubski, W. Infectious Etiologies of Parkinsonism: Pathomechanisms and Clinical Implications. *Front. Neurol.* **2019**, *10*, 652. [\[CrossRef\]](#)
92. Hawkes, C.H.; Del Tredici, K.; Braak, H. Parkinson's disease: A dual-hit hypothesis. *Neuropathol. Appl. Neurobiol.* **2007**, *33*, 599–614. [\[CrossRef\]](#)
93. Chen, Z.; Trapp, B.D. Microglia and neuroprotection. *J. Neurochem.* **2016**, *136*, 10–17. [\[CrossRef\]](#)
94. Tozzi, A.; Sciacaluga, M.; Loffredo, V.; Megaro, A.; Ledonne, A.; Cardinale, A.; Federici, M.; Bellingacci, L.; Paciotti, S.; Ferrari, E.; et al. Dopamine-dependent early synaptic and motor dysfunctions induced by α -synuclein in the nigrostriatal circuit. *Brain* **2021**, *144*, 3477–3491. [\[CrossRef\]](#) [\[PubMed\]](#)
95. Lehmann, J.M.; Lenhard, J.M.; Oliver, B.B.; Ringold, G.M.; Klierer, S.A. Peroxisome proliferator-activated receptors alpha and gamma are activated by indomethacin and other non-steroidal anti-inflammatory drugs. *J. Biol. Chem.* **1997**, *272*, 3406–3410. [\[CrossRef\]](#) [\[PubMed\]](#)
96. Kalampokini, S.; Becker, A.; Fassbender, K.; Lyros, E.; Unger, M.M. Nonpharmacological Modulation of Chronic Inflammation in Parkinson's Disease: Role of Diet Interventions. *Park. Dis.* **2019**, *2019*, e7535472. [\[CrossRef\]](#)
97. Zakiniaiez, Y.; Hoye, J.; Ryan Petrulli, J.; LeVasseur, B.; Stanley, G.; Gao, H.; Najafzadeh, S.; Ropchan, J.; Nabulsi, N.; Huang, Y.; et al. Systemic inflammation enhances stimulant-induced striatal dopamine elevation in tobacco smokers. *Brain Behav. Immun.* **2022**, *106*, 262–269. [\[CrossRef\]](#) [\[PubMed\]](#)
98. Moccia, M.; Picillo, M.; Erro, R.; Longo, K.; Amboni, M.; Santangelo, G.; Palladino, R.; Allocca, R.; Caporale, O.; Triassi, M.; et al. Increased bilirubin levels in de novo Parkinson's disease. *Eur. J. Neurol.* **2015**, *22*, 954–959. [\[CrossRef\]](#)
99. Sophiabadi, M.; Rastgoo, N.; Haghdost-Yazdi, H. Dopaminergic Neuronal Death in Substantia Nigra Associates with Serum Levels of Total Bilirubin, Selenium, and Zinc: Evidences from 6-Hydroxydopamine Animal Model of Parkinson's Disease. *Biol. Trace Elem. Res.* **2022**, *200*, 4058–4067. [\[CrossRef\]](#)
100. Hung, S.-Y.; Liou, H.-C.; Kang, K.-H.; Wu, R.-M.; Wen, C.-C.; Fu, W.-M. Over-expression of Heme oxygenase-1 protects dopaminergic neurons against 1-methyl-4-phenylpyridinium-induced neurotoxicity. *Mol. Pharmacol.* **2008**, *74*, 1564–1575. [\[CrossRef\]](#)
101. Yoo, M.S.; Chun, H.S.; Son, J.J.; DeGiorgio, L.A.; Kim, D.J.; Peng, C.; Son, J.H. Oxidative stress regulated genes in nigral dopaminergic neuronal cells: Correlation with the known pathology in Parkinson's disease. *Mol. Brain Res.* **2003**, *110*, 76–84. [\[CrossRef\]](#)
102. Ayuso, P.; Martínez, C.; Pastor, P.; Lorenzo-Betancor, O.; Luengo, A.; Jiménez-Jiménez, F.J.; Alonso-Navarro, H.; Agúndez, J.A.G.; García-Martín, E. An association study between Heme oxygenase-1 genetic variants and Parkinson's disease. *Front. Cell. Neurosci.* **2014**, *8*, 298. [\[CrossRef\]](#)
103. Schipper, H.M.; Song, W.; Zukor, H.; Hasclovici, J.R.; Zeligman, D. Heme oxygenase-1 and neurodegeneration: Expanding frontiers of engagement. *J. Neurochem.* **2009**, *110*, 469–485. [\[CrossRef\]](#)
104. Macías-García, D.; Méndez-Del Barrio, C.; Jesús, S.; Labrador, M.A.; Adarmes-Gómez, A.; Vargas-González, L.; Carrillo, F.; Gómez-Garre, P.; Mir, P. Increased bilirubin levels in Parkinson's disease. *Park. Relat. Disord.* **2019**, *63*, 213–216. [\[CrossRef\]](#)
105. Hatano, T.; Saiki, S.; Okuzumi, A.; Mohney, R.P.; Hattori, N. Identification of novel biomarkers for Parkinson's disease by metabolomic technologies. *J. Neurol. Neurosurg. Psychiatry* **2016**, *87*, 295–301. [\[CrossRef\]](#)
106. Kong, W.; Sun, X.; Yu, S.; Liu, P.; Zheng, X.; Zhang, J.; Zhu, L.; Jiang, T.; Jin, M.; Gao, J.; et al. Bile duct ligation increased dopamine levels in the cerebral cortex of rats partly due to induction of tyrosine hydroxylase. *Br. J. Pharmacol.* **2023**, *180*, 1690–1709. [\[CrossRef\]](#)
107. Berger, S.P.; Hüniger, M.; Yard, B.A.; Schnuelle, P.; Van Der Woude, F.J. Dopamine induces the expression of heme oxygenase-1 by human endothelial cells in vitro. *Kidney Int.* **2000**, *58*, 2314–2319. [\[CrossRef\]](#)
108. Pike, A.F.; Longhena, F.; Faustini, G.; van Eik, J.-M.; Gombert, I.; Herrebout, M.A.C.; Fayed, M.M.H.E.; Sandre, M.; Varanita, T.; Teunissen, C.E.; et al. Dopamine signaling modulates microglial NLRP3 inflammasome activation: Implications for Parkinson's disease. *J. Neuroinflammation* **2022**, *19*, 50. [\[CrossRef\]](#) [\[PubMed\]](#)
109. Amato, M.M.; Kilguss, N.V.; Gelardi, N.L.; Cashore, W.J. Dose-effect relationship of bilirubin on striatal synaptosomes in rats. *Biol. Neonate* **1994**, *66*, 288–293. [\[CrossRef\]](#)
110. Ochoa, E.L.; Wennberg, R.P.; An, Y.; Tandon, T.; Takashima, T.; Nguyen, T.; Chui, A. Interactions of bilirubin with isolated presynaptic nerve terminals: Functional effects on the uptake and release of neurotransmitters. *Cell. Mol. Neurobiol.* **1993**, *13*, 69–86. [\[CrossRef\]](#)
111. Brann, B.S.; Cashore, W.J.; Patrick, R.; Oh, W. in Vitro Effect of Bilirubin on Dopamine Synthesis in Adult Rat Brain Synaptosomes. *Pediatr. Res.* **1985**, *19*, 335. [\[CrossRef\]](#)
112. Roseth, S.; Hansen, T.W.R.; Fonnum, F.; Walaas, S.I. Bilirubin Inhibits Transport of Neurotransmitters in Synaptic Vesicles. *Pediatr. Res.* **1998**, *44*, 312–316. [\[CrossRef\]](#)

113. Effective Health Care Program, Agency for Healthcare Research and Quality, Rockville, MD. Chronic Idiopathic Hyperbilirubinemia and Parkinson's Disease. Content Last Reviewed February 2020. Available online: <https://effectivehealthcare.ahrq.gov/get-involved/nominated-topics/chronic-idiopathic-hyperbilirubinemia> (accessed on 27 June 2023).
114. Swenson, R.M.; Jew, J.Y. Learning deficits and brain monoamines in rats with congenital hyperbilirubinemia. *Exp. Neurol.* **1982**, *76*, 447–456. [\[CrossRef\]](#) [\[PubMed\]](#)
115. Miura, S.; Tsuchie, K.; Fukushima, M.; Arauchi, R.; Tsumori, T.; Otsuki, K.; Hayashida, M.; Hashioka, S.; Wake, R.; Miyaoka, T.; et al. Normalizing hyperactivity of the Gunn rat with bilirubin-induced neurological disorders via ketanserin. *Pediatr. Res.* **2022**, *91*, 556–564. [\[CrossRef\]](#)
116. Jayanti, S.; Moretti, R.; Tiribelli, C.; Gazzin, S. Bilirubin Prevents the TH+ Dopaminergic Neuron Loss in a Parkinson's Disease Model by Acting on TNF- α . *Int. J. Mol. Sci.* **2022**, *23*, 14276. [\[CrossRef\]](#) [\[PubMed\]](#)
117. Doorn, J.A.; Florang, V.R.; Schamp, J.H.; Vanle, B.C. Aldehyde dehydrogenase inhibition generates a reactive dopamine metabolite autotoxic to dopamine neurons. *Park. Relat. Disord.* **2014**, *20*, S73–S75. [\[CrossRef\]](#) [\[PubMed\]](#)
118. Costa-Mallen, P.; Gatenby, C.; Friend, S.; Maravilla, K.R.; Hu, S.-C.; Cain, K.C.; Agarwal, P.; Anzai, Y. Brain iron concentrations in regions of interest and relation with serum iron levels in Parkinson disease. *J. Neurol. Sci.* **2017**, *378*, 38–44. [\[CrossRef\]](#)
119. Hare, D.J.; Double, K.L. Iron and dopamine: A toxic couple. *Brain J. Neurol.* **2016**, *139 Pt 4*, 1026–1035. [\[CrossRef\]](#)
120. Song, L.; Song, W.; Schipper, H.M. Astroglia overexpressing heme oxygenase-1 predispose co-cultured PC12 cells to oxidative injury. *J. Neurosci. Res.* **2007**, *85*, 2186–2195. [\[CrossRef\]](#)
121. Dusek, P.; Hofer, T.; Alexander, J.; Roos, P.M.; Aaseth, J.O. Cerebral Iron Deposition in Neurodegeneration. *Biomolecules* **2022**, *12*, 714. [\[CrossRef\]](#) [\[PubMed\]](#)
122. Dusek, P.; Schneider, S.A. Neurodegeneration with brain iron accumulation. *Curr. Opin. Neurol.* **2012**, *25*, 499–506. [\[CrossRef\]](#)
123. Ndayisaba, A.; Kaindlstorfer, C.; Wenning, G.K. Iron in Neurodegeneration—Cause or Consequence? *Front. Neurosci.* **2019**, *13*, 180. [\[CrossRef\]](#)
124. Ward, R.J.; Dexter, D.T.; Crichton, R.R. Iron, Neuroinflammation and Neurodegeneration. *Int. J. Mol. Sci.* **2022**, *23*, 7267. [\[CrossRef\]](#)
125. Ward, R.J.; Zucca, F.A.; Duyn, J.H.; Crichton, R.R.; Zecca, L. The role of iron in brain ageing and neurodegenerative disorders. *Lancet Neurol.* **2014**, *13*, 1045–1060. [\[CrossRef\]](#) [\[PubMed\]](#)
126. Uranga, R.M.; Salvador, G.A. Unraveling the Burden of Iron in Neurodegeneration: Intersections with Amyloid Beta Peptide Pathology. *Oxid. Med. Cell. Longev.* **2018**, *2018*, 2850341. [\[CrossRef\]](#) [\[PubMed\]](#)
127. Marder, S.R.; Cannon, T.D. Schizophrenia. *N. Engl. J. Med.* **2019**, *381*, 1753–1761. [\[CrossRef\]](#)
128. McCutcheon, R.; Beck, K.; Jauhar, S.; Howes, O.D. Defining the Locus of Dopaminergic Dysfunction in Schizophrenia: A Meta-analysis and Test of the Mesolimbic Hypothesis. *Schizophr. Bull.* **2018**, *44*, 1301–1311. [\[CrossRef\]](#)
129. Benjamin, K.J.M.; Chen, Q.; Jaffe, A.E.; Stolz, J.M.; Collado-Torres, L.; Huuki-Myers, L.A.; Burke, E.E.; Arora, R.; Feltrin, A.S.; Barbosa, A.R.; et al. Analysis of the caudate nucleus transcriptome in individuals with schizophrenia highlights effects of antipsychotics and new risk genes. *Nat. Neurosci.* **2022**, *25*, 1559–1568. [\[CrossRef\]](#)
130. Weinberger, D.R. It's Dopamine and Schizophrenia All Over Again. *Biol. Psychiatry* **2022**, *92*, 757–759. [\[CrossRef\]](#) [\[PubMed\]](#)
131. Waddington, J.L. Psychosis in Parkinson's disease and parkinsonism in antipsychotic-naïve schizophrenia spectrum psychosis: Clinical, nosological and pathobiological challenges. *Acta Pharmacol. Sin.* **2020**, *41*, 464–470. [\[CrossRef\]](#)
132. Jaitpal, V.; Gawande, S. Olanzapine-Induced Parkinsonism and Akathisia: A Case Report. *Cureus* **2022**, *14*, e21354. [\[CrossRef\]](#) [\[PubMed\]](#)
133. Habermeyer, B.; Kneifel, S.; Lotz-Bläuer, I.; Müller-Spahn, F. Psychosis in a Case of Schizophrenia and Parkinson's Disease. *J. Neuropsychiatry Clin. Neurosci.* **2008**, *20*, 373–375. [\[CrossRef\]](#)
134. Ramzan, A.; Aggarwal, R.; Jamal, F. Novel case of Parkinson's disease and schizophrenia: Challenges in the management. *BMJ Case Rep. CP* **2021**, *14*, e244028. [\[CrossRef\]](#)
135. Gadit, A. Schizophrenia and Parkinson's disease: Challenges in management. *Case Rep.* **2011**, *2011*, bcr1120115108. [\[CrossRef\]](#) [\[PubMed\]](#)
136. Brown, A.S.; Derkits, E.J. Prenatal infection and schizophrenia: A review of epidemiologic and translational studies. *Am. J. Psychiatry* **2010**, *167*, 261–280. [\[CrossRef\]](#) [\[PubMed\]](#)
137. Khandaker, G.M.; Cousins, L.; Deakin, J.; Lennox, B.R.; Yolken, R.; Jones, P.B. Inflammation and immunity in schizophrenia: Implications for pathophysiology and treatment. *Lancet Psychiatry* **2015**, *2*, 258–270. [\[CrossRef\]](#) [\[PubMed\]](#)
138. Brown, A.S. Exposure to Prenatal Infection and Risk of Schizophrenia. *Front. Psychiatry* **2011**, *2*, 63. [\[CrossRef\]](#)
139. Jonakait, G.M. The effects of maternal inflammation on neuronal development: Possible mechanisms. *Int. J. Dev. Neurosci. Off. J. Int. Soc. Dev. Neurosci.* **2007**, *25*, 415–425. [\[CrossRef\]](#)
140. Patterson, P.H. Immune involvement in schizophrenia and autism: Etiology, pathology and animal models. *Behav. Brain Res.* **2009**, *204*, 313–321. [\[CrossRef\]](#)
141. Li, S.; Zhuo, M.; Huang, X.; Huang, Y.; Zhou, J.; Xiong, D.; Li, J.; Liu, Y.; Pan, Z.; Li, H.; et al. Altered gut microbiota associated with symptom severity in schizophrenia. *PeerJ* **2020**, *8*, e9574. [\[CrossRef\]](#)
142. Lin, C.; Chen, K.; Yu, J.; Feng, W.; Fu, W.; Yang, F.; Zhang, X.; Chen, D. Relationship between TNF- α levels and psychiatric symptoms in first-episode drug-naïve patients with schizophrenia before and after risperidone treatment and in chronic patients. *BMC Psychiatry* **2021**, *21*, 561. [\[CrossRef\]](#)

143. Lv, M.H.; Tan, Y.L.; Yan, S.X.; Tian, L.; Chen, D.C.; Tan, S.P.; Wang, Z.R.; Yang, F.D.; Yoon, J.H.; Zunta-Soares, G.B.; et al. Decreased serum TNF-alpha levels in chronic schizophrenia patients on long-term antipsychotics: Correlation with psychopathology and cognition. *Psychopharmacology* **2015**, *232*, 165–172. [[CrossRef](#)]
144. Hong, J.; Bang, M. Anti-inflammatory Strategies for Schizophrenia: A Review of Evidence for Therapeutic Applications and Drug Repurposing. *Clin. Psychopharmacol. Neurosci.* **2020**, *18*, 10–24. [[CrossRef](#)]
145. Albracht-Schulte, K.; Kalupahana, N.S.; Ramalingam, L.; Wang, S.; Rahman, S.M.; Robert-McComb, J.; Moustaid-Moussa, N. Omega-3 fatty acids in obesity and metabolic syndrome: A mechanistic update. *J. Nutr. Biochem.* **2018**, *58*, 1–16. [[CrossRef](#)] [[PubMed](#)]
146. Cha, H.Y.; Yang, S.J. Anti-Inflammatory Diets and Schizophrenia. *Clin. Nutr. Res.* **2020**, *9*, 241–257. [[CrossRef](#)]
147. Becklén, M.; Orhan, F.; Piehl, F.; Cervenka, S.; Sellgren, C.M.; Flyckt, L.; Erhardt, S.; Fatouros-Bergman, H. Plasma bilirubin levels are reduced in first-episode psychosis patients and associates to working memory and duration of untreated psychosis. *Sci. Rep.* **2021**, *11*, 7527. [[CrossRef](#)]
148. Radhakrishnan, R.; Kanigere, M.; Menon, J.; Calvin, S.; Janish, A.; Srinivasan, K. Association between unconjugated bilirubin and schizophrenia. *Psychiatry Res.* **2011**, *189*, 480–482. [[CrossRef](#)] [[PubMed](#)]
149. Miyaoka, T. Schizophrenia-Associated Idiopathic Unconjugated Hyperbilirubinemia (Gilbert's Syndrome). *J. Clin. Psychiatry* **2000**, *61*, 868–871. [[CrossRef](#)]
150. Uutela, P.; Karhu, L.; Piepponen, P.; Käenmäki, M.; Ketola, R.A.; Kostiainen, R. Discovery of Dopamine Glucuronide in Rat and Mouse Brain Microdialysis Samples Using Liquid Chromatography Tandem Mass Spectrometry. *Anal. Chem.* **2009**, *81*, 427–434. [[CrossRef](#)] [[PubMed](#)]
151. Llido, J.P.; Fioriti, E.; Pascut, D.; Giuffrè, M.; Bottin, C.; Zanconati, F.; Tiribelli, C.; Gazzin, S. Bilirubin-Induced Transcriptomic Imprinting in Neonatal Hyperbilirubinemia. *Biology* **2023**, *12*, 834. [[CrossRef](#)]
152. Dornelles, E.P.; Marques, J.G.; Ouakinin, S. Unconjugated bilirubin and schizophrenia: A systematic review. *CNS Spectr.* **2011**, *189*, 480–482. [[CrossRef](#)]
153. Gazzin, S.; Jayanti, S.; Tiribelli, C. Models of bilirubin neurological damage: Lessons learned and new challenges. *Pediatr. Res.* **2022**, *1*–8. [[CrossRef](#)]
154. Song, W.; Zukor, H.; Lin, S.H.; Hascalovici, J.; Liberman, A.; Tavitian, A.; Mui, J.; Vali, H.; Tong, X.-K.; Bhardwaj, S.; et al. Schizophrenia-Like Features in Transgenic Mice Overexpressing Human HO-1 in the Astrocytic Compartment. *J. Neurosci. Off. J. Soc. Neurosci.* **2012**, *32*, 10841–10853. [[CrossRef](#)]
155. Drechsler, R.; Brem, S.; Brandeis, D.; Grünblatt, E.; Berger, G.; Walitza, S. ADHD: Current Concepts and Treatments in Children and Adolescents. *Neuropediatrics* **2020**, *51*, 315–335. [[CrossRef](#)] [[PubMed](#)]
156. Klein, M.O.; Battagello, D.S.; Cardoso, A.R.; Hauser, D.N.; Bittencourt, J.C.; Correa, R.G. Dopamine: Functions, Signaling, and Association with Neurological Diseases. *Cell. Mol. Neurobiol.* **2019**, *39*, 31–59. [[CrossRef](#)] [[PubMed](#)]
157. Cortese, S.; Coghill, D. Twenty years of research on attention-deficit/hyperactivity disorder (ADHD): Looking back, looking forward. *BMJ Ment Health* **2018**, *21*, 173–176. [[CrossRef](#)] [[PubMed](#)]
158. Han, V.X.; Patel, S.; Jones, H.F.; Dale, R.C. Maternal immune activation and neuroinflammation in human neurodevelopmental disorders. *Nat. Rev. Neurol.* **2021**, *17*, 564–579. [[CrossRef](#)]
159. Becker, K.; El-Faddagh, M.; Schmidt, M.H.; Esser, G.; Laucht, M. Interaction of dopamine transporter genotype with prenatal smoke exposure on ADHD symptoms. *J. Pediatr.* **2008**, *152*, 263–269. [[CrossRef](#)]
160. Nigg, J.; Nikolas, M.; Burt, S.A. Measured gene-by-environment interaction in relation to attention-deficit/hyperactivity disorder. *J. Am. Acad. Child Adolesc. Psychiatry* **2010**, *49*, 863–873. [[CrossRef](#)]
161. Wu, J.; Xiao, H.; Sun, H.; Zou, L.; Zhu, L.-Q. Role of Dopamine Receptors in ADHD: A Systematic Meta-analysis. *Mol. Neurobiol.* **2012**, *45*, 605–620. [[CrossRef](#)]
162. Czerniak, S.M.; Sikoglu, E.M.; King, J.A.; Kennedy, D.N.; Mick, E.; Frazier, J.; Moore, C.M. Areas of the brain modulated by single-dose methylphenidate treatment in youth with ADHD during task-based fMRI: A systematic review. *Harv. Rev. Psychiatry* **2013**, *21*, 151–162. [[CrossRef](#)]
163. Faraone, S.V.; Perlis, R.H.; Doyle, A.E.; Smoller, J.W.; Goralnick, J.J.; Holmgren, M.A.; Sklar, P. Molecular Genetics of Attention-Deficit/Hyperactivity Disorder. *Biol. Psychiatry* **2005**, *57*, 1313–1323. [[CrossRef](#)]
164. Gilbert, D.L.; Wang, Z.; Sallee, F.R.; Ridel, K.R.; Merhar, S.; Zhang, J.; Lipps, T.D.; White, C.; Badreldin, N.; Wassermann, E.M. Dopamine transporter genotype influences the physiological response to medication in ADHD. *Brain* **2006**, *129*, 2038–2046. [[CrossRef](#)]
165. Volkow, N.D.; Wang, G.-J.; Newcorn, J.; Fowler, J.S.; Telang, F.; Solanto, M.V.; Logan, J.; Wong, C.; Ma, Y.; Swanson, J.M.; et al. Brain dopamine transporter levels in treatment and drug naïve adults with ADHD. *NeuroImage* **2007**, *34*, 1182–1190. [[CrossRef](#)]
166. Del Campo, N.; Chamberlain, S.R.; Sahakian, B.J.; Robbins, T.W. The roles of dopamine and noradrenaline in the pathophysiology and treatment of attention-deficit/hyperactivity disorder. *Biol. Psychiatry* **2011**, *69*, e145–e157. [[CrossRef](#)] [[PubMed](#)]
167. Dunn, G.A.; Nigg, J.T.; Sullivan, E.L. Neuroinflammation as a risk factor for attention deficit hyperactivity disorder. *Pharmacol. Biochem. Behav.* **2019**, *182*, 22–34. [[CrossRef](#)] [[PubMed](#)]
168. Anand, D.; Colpo, G.D.; Zeni, G.; Zeni, C.P.; Teixeira, A.L. Attention-Deficit/Hyperactivity Disorder And Inflammation: What Does Current Knowledge Tell Us? A Systematic Review. *Front. Psychiatry* **2017**, *8*, 228. [[CrossRef](#)] [[PubMed](#)]

169. Saccaro, L.F.; Schilliger, Z.; Perroud, N.; Piguet, C. Inflammation, Anxiety, and Stress in Attention-Deficit/Hyperactivity Disorder. *Biomedicines* **2021**, *9*, 1313. [\[CrossRef\]](#)
170. Leffa, D.T.; Torres, I.L.S.; Rohde, L.A. A Review on the Role of Inflammation in Attention-Deficit/Hyperactivity Disorder. *Neuroimmunomodulation* **2018**, *25*, 328–333. [\[CrossRef\]](#)
171. Park, J.H. Potential Inflammatory Biomarker in Patients with Attention Deficit Hyperactivity Disorder. *Int. J. Mol. Sci.* **2022**, *23*, 13054. [\[CrossRef\]](#)
172. O'Shea, T.M.; Joseph, R.M.; Kuban, K.C.K.; Allred, E.N.; Ware, J.; Coster, T.; Fichorova, R.N.; Dammann, O.; Leviton, A. Elevated blood levels of inflammation-related proteins are associated with an attention problem at age 24 mo in extremely preterm infants. *Pediatr. Res.* **2014**, *75*, 781–787. [\[CrossRef\]](#)
173. Cortese, S.; Angriman, M.; Comencini, E.; Vincenzi, B.; Maffei, C. Association between inflammatory cytokines and ADHD symptoms in children and adolescents with obesity: A pilot study. *Psychiatry Res.* **2019**, *278*, 7–11. [\[CrossRef\]](#)
174. Demontis, D.; Walters, R.K.; Martin, J.; Mattheisen, M.; Als, T.D.; Agerbo, E.; Baldursson, G.; Belliveau, R.; Bybjerg-Grauholm, J.; Bækvad-Hansen, M.; et al. Discovery of the first genome-wide significant risk loci for attention deficit/hyperactivity disorder. *Nat. Genet.* **2019**, *51*, 63–75. [\[CrossRef\]](#)
175. Enard, W.; Gehre, S.; Hammerschmidt, K.; Hölter, S.M.; Blass, T.; Somel, M.; Brückner, M.K.; Schreiweis, C.; Winter, C.; Sohr, R.; et al. A humanized version of Foxp2 affects cortico-basal ganglia circuits in mice. *Cell* **2009**, *137*, 961–971. [\[CrossRef\]](#) [\[PubMed\]](#)
176. Hoogman, M.; Bralten, J.; Hibar, D.P.; Mennes, M.; Zwiers, M.P.; Schweren, L.S.J.; van Hulzen, K.J.E.; Medland, S.E.; Shumskaya, E.; Jahanshad, N.; et al. Subcortical brain volume differences in participants with attention deficit hyperactivity disorder in children and adults: A cross-sectional mega-analysis. *Lancet Psychiatry* **2017**, *4*, 310–319. [\[CrossRef\]](#) [\[PubMed\]](#)
177. Wei, C.-C.; Chang, C.-H.; Lin, C.-L.; Chang, S.-N.; Li, T.-C.; Kao, C.-H. Neonatal jaundice and increased risk of attention-deficit hyperactivity disorder: A population-based cohort study. *J. Child Psychol. Psychiatry* **2015**, *56*, 460–467. [\[CrossRef\]](#) [\[PubMed\]](#)
178. Hokkanen, L.; Launes, J.; Michelsson, K. Adult neurobehavioral outcome of hyperbilirubinemia in full term neonates—A 30 year prospective follow-up study. *PeerJ* **2014**, *2*, e294. [\[CrossRef\]](#)
179. Amin, S.B.; Smith, T.; Timler, G. Developmental influence of unconjugated hyperbilirubinemia and neurobehavioral disorders. *Pediatr. Res.* **2019**, *85*, 191–197. [\[CrossRef\]](#)
180. Vázquez, J.C.; Martín de la Torre, O.; López Palomé, J.; Redolar-Ripoll, D. Effects of Caffeine Consumption on Attention Deficit Hyperactivity Disorder (ADHD) Treatment: A Systematic Review of Animal Studies. *Nutrients* **2022**, *14*, 739. [\[CrossRef\]](#)
181. Zhou, W.-Z.; Ye, A.Y.; Sun, Z.-K.; Tian, H.H.; Pu, T.Z.; Wu, Y.-Y.; Wang, D.-D.; Zhao, M.-Z.; Lu, S.-J.; Yang, C.-H.; et al. Statistical analysis of twenty years (1993 to 2012) of data from mainland China's first intervention center for children with autism spectrum disorder. *Mol. Autism* **2014**, *5*, 52. [\[CrossRef\]](#)
182. Ernst, M.; Zametkin, A.J.; Matochik, J.A.; Pascualvaca, D.; Cohen, R.M. Low medial prefrontal dopaminergic activity in autistic children. *Lancet Lond. Engl.* **1997**, *350*, 638. [\[CrossRef\]](#)
183. Chevallier, C.; Kohls, G.; Troiani, V.; Brodtkin, E.S.; Schultz, R.T. The social motivation theory of autism. *Trends Cogn. Sci.* **2012**, *16*, 231–239. [\[CrossRef\]](#)
184. Masi, A.; Quintana, D.S.; Glozier, N.; Lloyd, A.R.; Hickie, I.B.; Guastella, A.J. Cytokine aberrations in autism spectrum disorder: A systematic review and meta-analysis. *Mol. Psychiatry* **2015**, *20*, 440–446. [\[CrossRef\]](#)
185. Ferguson, B.J.; Marler, S.; Altstein, L.L.; Lee, E.B.; Mazurek, M.O.; McLaughlin, A.; Macklin, E.A.; McDonnell, E.; Davis, D.J.; Belenchia, A.M.; et al. Associations between cytokines, endocrine stress response, and gastrointestinal symptoms in autism spectrum disorder. *Brain. Behav. Immun.* **2016**, *58*, 57–62. [\[CrossRef\]](#) [\[PubMed\]](#)
186. Krakowiak, P.; Goines, P.E.; Tancredi, D.J.; Ashwood, P.; Hansen, R.L.; Hertz-Picciotto, I.; Van de Water, J. Neonatal Cytokine Profiles Associated With Autism Spectrum Disorder. *Biol. Psychiatry* **2017**, *81*, 442–451. [\[CrossRef\]](#)
187. Xu, N.; Li, X.; Zhong, Y. Inflammatory cytokines: Potential biomarkers of immunologic dysfunction in autism spectrum disorders. *Mediators Inflamm.* **2015**, *2015*, 531518. [\[CrossRef\]](#) [\[PubMed\]](#)
188. Xie, J.; Huang, L.; Li, X.; Li, H.; Zhou, Y.; Zhu, H.; Pan, T.; Kendrick, K.M.; Xu, W. Immunological cytokine profiling identifies TNF- α as a key molecule dysregulated in autistic children. *Oncotarget* **2017**, *8*, 82390–82398. [\[CrossRef\]](#) [\[PubMed\]](#)
189. Jenabi, E.; Bashirian, S.; Khazaei, S. Association between neonatal jaundice and autism spectrum disorders among children: A meta-analysis. *Clin. Exp. Pediatr.* **2020**, *63*, 8–13. [\[CrossRef\]](#)
190. Hao, Y.; Ge, H.; Sun, M.; Gao, Y. Selecting an Appropriate Animal Model of Depression. *Int. J. Mol. Sci.* **2019**, *20*, 4827. [\[CrossRef\]](#)
191. Tye, K.M.; Mirzabekov, J.J.; Warden, M.R.; Ferenczi, E.A.; Tsai, H.-C.; Finkelstein, J.; Kim, S.-Y.; Adhikari, A.; Thompson, K.R.; Andalman, A.S.; et al. Dopamine neurons modulate neural encoding and expression of depression-related behaviour. *Nature* **2013**, *493*, 537–541. [\[CrossRef\]](#)
192. Beurel, E.; Toups, M.; Nemeroff, C.B. The Bidirectional Relationship of Depression and Inflammation: Double Trouble. *Neuron* **2020**, *107*, 234–256. [\[CrossRef\]](#) [\[PubMed\]](#)
193. Lucido, M.J.; Bekhbat, M.; Goldsmith, D.R.; Treadway, M.T.; Haroon, E.; Felger, J.C.; Miller, A.H. Aiding and Abetting Anhedonia: Impact of Inflammation on the Brain and Pharmacological Implications. *Pharmacol. Rev.* **2021**, *73*, 1084–1117. [\[CrossRef\]](#)
194. Goldsmith, D.R.; Bekhbat, M.; Le, N.-A.; Chen, X.; Woolwine, B.J.; Li, Z.; Haroon, E.; Felger, J.C. Protein and gene markers of metabolic dysfunction and inflammation together associate with functional connectivity in reward and motor circuits in depression. *Brain. Behav. Immun.* **2020**, *88*, 193–202. [\[CrossRef\]](#)

195. Jeon, S.W.; Kim, Y.K. Neuroinflammation and cytokine abnormality in major depression: Cause or consequence in that illness? *World J. Psychiatry* **2016**, *6*, 283–293. [[CrossRef](#)]
196. Cheng, Y.; Desse, S.; Martinez, A.; Worthen, R.J.; Jope, R.S.; Beurel, E. TNF α disrupts blood brain barrier integrity to maintain prolonged depressive-like behavior in mice. *Brain. Behav. Immun.* **2018**, *69*, 556–567. [[CrossRef](#)] [[PubMed](#)]
197. Morgan, J.A.; Singhal, G.; Corrigan, F.; Jaehne, E.J.; Jawahar, M.C.; Baune, B.T. Exercise related anxiety-like behaviours are mediated by TNF receptor signaling, but not depression-like behaviours. *Brain Res.* **2018**, *1695*, 10–17. [[CrossRef](#)]
198. Raison, C.L.; Rutherford, R.E.; Woolwine, B.J.; Shuo, C.; Schettler, P.; Drake, D.F.; Haroon, E.; Miller, A.H. A randomized controlled trial of the tumor necrosis factor antagonist infliximab for treatment-resistant depression: The role of baseline inflammatory biomarkers. *JAMA Psychiatry* **2013**, *70*, 31–41. [[CrossRef](#)] [[PubMed](#)]
199. Müller, N.; Schwarz, M.J.; Dehning, S.; Douhe, A.; Ceroveck, A.; Goldstein-Müller, B.; Spellmann, I.; Hetzel, G.; Maino, K.; Kleindienst, N.; et al. The cyclooxygenase-2 inhibitor celecoxib has therapeutic effects in major depression: Results of a double-blind, randomized, placebo controlled, add-on pilot study to reboxetine. *Mol. Psychiatry* **2006**, *11*, 680–684. [[CrossRef](#)] [[PubMed](#)]
200. Kohno, M.; Link, J.; Dennis, L.E.; McCready, H.; Huckans, M.; Hoffman, W.F.; Loftis, J.M. Neuroinflammation in addiction: A review of neuroimaging studies and potential immunotherapies. *Pharmacol. Biochem. Behav.* **2019**, *179*, 34–42. [[CrossRef](#)] [[PubMed](#)]
201. Peng, Y.-F.; Xiang, Y.; Wei, Y.-S. The significance of routine biochemical markers in patients with major depressive disorder. *Sci. Rep.* **2016**, *6*, 34402. [[CrossRef](#)]
202. Oren, D.A.; Desan, P.H.; Boutros, N.; Anand, A.; Charney, D.S. Effects of light on low nocturnal bilirubin in winter depression: A preliminary report. *Biol. Psychiatry* **2002**, *51*, 422–425. [[CrossRef](#)]
203. Wise, R.A.; Robble, M.A. Dopamine and Addiction. *Annu. Rev. Psychol.* **2020**, *71*, 79–106. [[CrossRef](#)]
204. Scofield, M.D.; Heinsbroek, J.A.; Gipson, C.D.; Kupchik, Y.M.; Spencer, S.; Smith, A.C.W.; Roberts-Wolfe, D.; Kalivas, P.W. The Nucleus Accumbens: Mechanisms of Addiction across Drug Classes Reflect the Importance of Glutamate Homeostasis. *Pharmacol. Rev.* **2016**, *68*, 816–871. [[CrossRef](#)]
205. Broom, S.L.; Yamamoto, B.K. Effects of subchronic methamphetamine exposure on basal dopamine and stress-induced dopamine release in the nucleus accumbens shell of rats. *Psychopharmacology* **2005**, *181*, 467–476. [[CrossRef](#)] [[PubMed](#)]
206. Baik, J.-H. Dopamine signaling in food addiction: Role of dopamine D2 receptors. *BMB Rep.* **2013**, *46*, 519–526. [[CrossRef](#)]
207. Wang, G.J.; Volkow, N.D.; Logan, J.; Pappas, N.R.; Wong, C.T.; Zhu, W.; Netusil, N.; Fowler, J.S. Brain dopamine and obesity. *Lancet* **2001**, *357*, 354–357. [[CrossRef](#)] [[PubMed](#)]
208. Wallace, C.W.; Fordahl, S.C. Obesity and dietary fat influence dopamine neurotransmission: Exploring the convergence of metabolic state, physiological stress, and inflammation on dopaminergic control of food intake. *Nutr. Res. Rev.* **2022**, *35*, 236–251. [[CrossRef](#)] [[PubMed](#)]
209. MA, H.; ZHU, G. The dopamine system and alcohol dependence. *Shanghai Arch. Psychiatry* **2014**, *26*, 61–68. [[CrossRef](#)] [[PubMed](#)]
210. Nguyen, C.; Mondoloni, S.; Le Borgne, T.; Centeno, I.; Come, M.; Jehl, J.; Solié, C.; Reynolds, L.M.; Durand-de Cuttoli, R.; Tolu, S.; et al. Nicotine inhibits the VTA-to-amygdala dopamine pathway to promote anxiety. *Neuron* **2021**, *109*, 2604–2615.e9. [[CrossRef](#)] [[PubMed](#)]
211. Ransohoff, R.M. Microgliosis: The questions shape the answers. *Nat. Neurosci.* **2007**, *10*, 1507–1509. [[CrossRef](#)] [[PubMed](#)]
212. Hanisch, U.-K. Microglia as a source and target of cytokines. *Glia* **2002**, *40*, 140–155. [[CrossRef](#)]
213. Otten, U.; März, P.; Heese, K.; Hock, C.; Kunz, D.; Rose-John, S. Cytokines and neurotrophins interact in normal and diseased states. *Ann. N. Y. Acad. Sci.* **2000**, *917*, 322–330. [[CrossRef](#)]
214. Beardsley, P.M.; Hauser, K.F. Chapter One—Glial Modulators as Potential Treatments of Psychostimulant Abuse. In *Advances in Pharmacology*; Dwoskin, L.P., Ed.; Emerging Targets & Therapeutics in the Treatment of Psychostimulant Abuse; Academic Press: Cambridge, MA, USA, 2014; Volume 69, pp. 1–69.
215. Maeda, T.; Kiguchi, N.; Fukazawa, Y.; Yamamoto, A.; Ozaki, M.; Kishioka, S. Peroxisome Proliferator-Activated Receptor Gamma Activation Relieves Expression of Behavioral Sensitization to Methamphetamine in Mice. *Neuropsychopharmacology* **2007**, *32*, 1133–1140. [[CrossRef](#)]
216. Nakajima, A.; Yamada, K.; Nagai, T.; Uchiyama, T.; Miyamoto, Y.; Mamiya, T.; He, J.; Nitta, A.; Mizuno, M.; Tran, M.H.; et al. Role of tumor necrosis factor-alpha in methamphetamine-induced drug dependence and neurotoxicity. *J. Neurosci. Off. J. Soc. Neurosci.* **2004**, *24*, 2212–2225. [[CrossRef](#)] [[PubMed](#)]
217. Lewitus, G.M.; Konefal, S.C.; Greenhalgh, A.D.; Pribrag, H.; Augereau, K.; Stellwagen, D. Microglial TNF- α Suppresses Cocaine-Induced Plasticity and Behavioral Sensitization. *Neuron* **2016**, *90*, 483–491. [[CrossRef](#)]
218. Rizzo, M.D.; Crawford, R.B.; Henriquez, J.E.; Aldhamen, Y.A.; Gulick, P.; Amalfitano, A.; Kaminski, N.E. HIV-infected cannabis users have lower circulating CD16+ monocytes and IFN- γ -inducible protein 10 levels compared with nonusing HIV patients. *AIDS Lond. Engl.* **2018**, *32*, 419–429. [[CrossRef](#)] [[PubMed](#)]
219. Bayazit, H.; Selek, S.; Karababa, I.F.; Cicek, E.; Aksoy, N. Evaluation of Oxidant/Antioxidant Status and Cytokine Levels in Patients with Cannabis Use Disorder. *Clin. Psychopharmacol. Neurosci. Off. Sci. J. Korean Coll. Neuropsychopharmacol.* **2017**, *15*, 237–242. [[CrossRef](#)] [[PubMed](#)]

220. Krey, G.; Braissant, O.; L'Horsset, F.; Kalkhoven, E.; Perroud, M.; Parker, M.G.; Wahli, W. Fatty acids, eicosanoids, and hypolipidemic agents identified as ligands of peroxisome proliferator-activated receptors by coactivator-dependent receptor ligand assay. *Mol. Endocrinol.* **1997**, *11*, 779–791. [\[CrossRef\]](#)
221. Chen, H.; Uz, T.; Manev, H. Minocycline affects cocaine sensitization in mice. *Neurosci. Lett.* **2009**, *452*, 258–261. [\[CrossRef\]](#)
222. Lee, H.-S.; Jeong, G.-S. 6,7,4'-Trihydroxyflavanone Mitigates Methamphetamine-Induced Neurotoxicity in SH-SY5y Cells via Nrf2/heme Oxygenase-1 and PI3K/Akt/mTOR Signaling Pathways. *Molecules* **2021**, *26*, 2442. [\[CrossRef\]](#)
223. Belayev, L.; Hong, S.-H.; Menghani, H.; Marcell, S.J.; Obenaus, A.; Freitas, R.S.; Khoutorova, L.; Balaszczuk, V.; Jun, B.; Oriá, R.B.; et al. Docosanoids Promote Neurogenesis and Angiogenesis, Blood-Brain Barrier Integrity, Penumbra Protection, and Neurobehavioral Recovery After Experimental Ischemic Stroke. *Mol. Neurobiol.* **2018**, *55*, 7090–7106. [\[CrossRef\]](#)
224. Hashimoto, N.; Matsumoto, T.; Mabe, H.; Hashitani, T.; Nishino, H. Dopamine has inhibitory and accelerating effects on ischemia-induced neuronal cell damage in the rat striatum. *Brain Res. Bull.* **1994**, *33*, 281–288. [\[CrossRef\]](#)
225. Akiyama, Y.; Ito, A.; Koshimura, K.; Ohue, T.; Yamagata, S.; Miwa, S.; Kikuchi, H. Effects of transient forebrain ischemia and reperfusion on function of dopaminergic neurons and dopamine reuptake in vivo in rat striatum. *Brain Res.* **1991**, *561*, 120–127. [\[CrossRef\]](#)
226. Sieber, M.W.; Guenther, M.; Jaenisch, N.; Albrecht-Eckardt, D.; Kohl, M.; Witte, O.W.; Frahm, C. Age-specific transcriptional response to stroke. *Neurobiol. Aging* **2014**, *35*, 1744–1754. [\[CrossRef\]](#) [\[PubMed\]](#)
227. Momosaki, S.; Ito, M.; Yamato, H.; Iimori, H.; Sumiyoshi, H.; Morimoto, K.; Imamoto, N.; Watabe, T.; Shimosegawa, E.; Hatazawa, J.; et al. Longitudinal imaging of the availability of dopamine transporter and D2 receptor in rat striatum following mild ischemia. *J. Cereb. Blood Flow Metab.* **2017**, *37*, 605–613. [\[CrossRef\]](#) [\[PubMed\]](#)
228. Lam, J.M.; Globas, C.; Hosp, J.A.; Karnath, H.-O.; Wächter, T.; Luft, A.R. Impaired implicit learning and feedback processing after stroke. *Neuroscience* **2016**, *314*, 116–124. [\[CrossRef\]](#) [\[PubMed\]](#)
229. Niewiarowska-Sendo, A.; Kozik, A.; Guevara-Lora, I. Influence of bradykinin B2 receptor and dopamine D2 receptor on the oxidative stress, inflammatory response, and apoptotic process in human endothelial cells. *PLoS ONE* **2018**, *13*, e0206443. [\[CrossRef\]](#)
230. Winkelman, J.W.; Armstrong, M.J.; Allen, R.P.; Chaudhuri, K.R.; Ondo, W.; Trenkwalder, C.; Zee, P.C.; Gronseth, G.S.; Gloss, D.; Zesiewicz, T. Practice guideline summary: Treatment of restless legs syndrome in adults: Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. *Neurology* **2016**, *87*, 2585–2593. [\[CrossRef\]](#)
231. Kaushik, P.; Ali, M.; Tabassum, H.; Parvez, S. Post-ischemic administration of dopamine D2 receptor agonist reduces cell death by activating mitochondrial pathway following ischemic stroke. *Life Sci.* **2020**, *261*, 118349. [\[CrossRef\]](#)
232. Wang, H.; Du, Y.; Xu, W.; Li, C.; Sun, H.; Hu, K.; Hu, Y.; Yu, T.; Guo, H.; Xie, L.; et al. Exogenous glutathione exerts a therapeutic effect in ischemic stroke rats by interacting with intrastriatal dopamine. *Acta Pharmacol. Sin.* **2022**, *43*, 541–551. [\[CrossRef\]](#)
233. Parikh, N.S.; Merkler, A.E.; Iadecola, C. Inflammation, Autoimmunity, Infection, and Stroke. *Stroke* **2020**, *51*, 711–718. [\[CrossRef\]](#)
234. Choi, Y.; Lee, S.J.; Spiller, W.; Jung, K.J.; Lee, J.-Y.; Kimm, H.; Back, J.H.; Lee, S.; Jee, S.H. Causal Associations Between Serum Bilirubin Levels and Decreased Stroke Risk: A Two-Sample Mendelian Randomization Study. *Arterioscler. Thromb. Vasc. Biol.* **2020**, *40*, 437–445. [\[CrossRef\]](#)
235. Zhong, K.; Wang, X.; Ma, X.; Ji, X.; Sang, S.; Shao, S.; Zhao, Y.; Xiang, Y.; Li, J.; Wang, G.; et al. Association between serum bilirubin and asymptomatic intracranial atherosclerosis: Results from a population-based study. *Neurol. Sci.* **2020**, *41*, 1531–1538. [\[CrossRef\]](#)
236. Targher, G. Risk of Ischemic Stroke and Decreased Serum Bilirubin Levels. *Arterioscler. Thromb. Vasc. Biol.* **2014**, *34*, 702–704. [\[CrossRef\]](#) [\[PubMed\]](#)
237. Stocker, R. Antioxidant Activities of Bile Pigments. *Antioxid. Redox Signal.* **2004**, *6*, 841–849. [\[CrossRef\]](#)
238. Stocker, R.; Yamamoto, Y.; McDonagh, A.F.; Glazer, A.N.; Ames, B.N. Bilirubin is an antioxidant of possible physiological importance. *Science* **1987**, *235*, 1043–1046. [\[CrossRef\]](#) [\[PubMed\]](#)
239. Baranano, D.E.; Rao, M.; Ferris, C.D.; Snyder, S.H. Biliverdin reductase: A major physiologic cytoprotectant. *Proc. Natl. Acad. Sci. USA* **2002**, *99*, 16093–16098. [\[CrossRef\]](#) [\[PubMed\]](#)
240. Melnikov, M.; Pashenkov, M.; Boyko, A. Dopaminergic Receptor Targeting in Multiple Sclerosis: Is There Therapeutic Potential? *Int. J. Mol. Sci.* **2021**, *22*, 5313. [\[CrossRef\]](#)
241. Carandini, T.; Mancini, M.; Bogdan, I.; Rae, C.L.; Barritt, A.W.; Sethi, A.; Harrison, N.; Rashid, W.; Scarpini, E.; Galimberti, D.; et al. Disruption of brainstem monoaminergic fibre tracts in multiple sclerosis as a putative mechanism for cognitive fatigue: A fixel-based analysis. *NeuroImage Clin.* **2021**, *30*, 102587. [\[CrossRef\]](#) [\[PubMed\]](#)
242. Levite, M.; Marino, F.; Cosentino, M. Dopamine, T cells and multiple sclerosis (MS). *J. Neural Transm.* **2017**, *124*, 525–542. [\[CrossRef\]](#)
243. Giorelli, M.; Livrea, P.; Trojano, M. Dopamine Fails to Regulate Activation of Peripheral Blood Lymphocytes from Multiple Sclerosis Patients: Effects of IFN- β . *J. Interferon Cytokine Res.* **2005**, *25*, 395–406. [\[CrossRef\]](#)
244. Alvarenga-Filho, H.; Salles, M.; Hygino, J.; Ferreira, T.B.; Sacramento, P.M.; Monteiro, C.; Vasconcelos, C.C.F.; Alvarenga, R.M.P.; Bento, C.A.M. Fatigue favors in vitro Th1 and Th17-like cell expansion and reduces corticoid sensitivity in MS patients. *J. Neuroimmunol.* **2017**, *303*, 81–89. [\[CrossRef\]](#)

245. Grzegorski, T.; Iwanowski, P.; Kozubski, W.; Losy, J. The alterations of cerebrospinal fluid TNF-alpha and TGF-beta2 levels in early relapsing-remitting multiple sclerosis. *Immunol. Res.* **2022**, *70*, 708–713. [CrossRef]
246. Dobryakova, E.; Genova, H.M.; DeLuca, J.; Wylie, G.R. The Dopamine Imbalance Hypothesis of Fatigue in Multiple Sclerosis and Other Neurological Disorders. *Front. Neurol.* **2015**, *6*, 52. [CrossRef]
247. Liu, Y.; Zhu, B.; Luo, L.; Li, P.; Paty, D.W.; Cynader, M.S. Heme oxygenase-1 plays an important protective role in experimental autoimmune encephalomyelitis. *Neuroreport* **2001**, *12*, 1841–1845. [CrossRef]
248. Liu, Y.; Liu, J.; Tetzlaff, W.; Paty, D.W.; Cynader, M.S. Biliverdin reductase, a major physiologic cytoprotectant, suppresses experimental autoimmune encephalomyelitis. *Free Radic. Biol. Med.* **2006**, *40*, 960–967. [CrossRef] [PubMed]
249. Liu, Y.; Zhu, B.; Wang, X.; Luo, L.; Li, P.; Paty, D.W.; Cynader, M.S. Bilirubin as a potent antioxidant suppresses experimental autoimmune encephalomyelitis: Implications for the role of oxidative stress in the development of multiple sclerosis. *J. Neuroimmunol.* **2003**, *139*, 27–35. [CrossRef] [PubMed]
250. Kim, T.W.; Kim, Y.; Jung, W.; Kim, D.E.; Keum, H.; Son, Y.; Jon, S. Bilirubin nanomedicine ameliorates the progression of experimental autoimmune encephalomyelitis by modulating dendritic cells. *J. Control. Release* **2021**, *331*, 74–84. [CrossRef]
251. Amin, S.B.; Smith, T.; Wang, H. Is Neonatal Jaundice Associated with Autism Spectrum Disorders: A Systematic Review. *J. Autism Dev. Disord.* **2011**, *41*, 1455–1463. [CrossRef] [PubMed]
252. Dopamine Treatment in Children With Cerebral Palsy With Dystonia—A Double Blind Controlled Study—Full Text View—ClinicalTrials.gov. Available online: <https://clinicaltrials.gov/ct2/show/NCT01361373> (accessed on 7 July 2023).
253. Paton, M.C.B.; Finch-Edmondson, M.; Dale, R.C.; Fahey, M.C.; Nold-Petry, C.A.; Nold, M.F.; Griffin, A.R.; Novak, I. Persistent Inflammation in Cerebral Palsy: Pathogenic Mediator or Comorbidity? A Scoping Review. *J. Clin. Med.* **2022**, *11*, 7368. [CrossRef] [PubMed]
254. Zareen, Z.; Strickland, T.; Fallah, L.; McEneaney, V.; Kelly, L.; McDonald, D.; Molloy, E.J. Cytokine dysregulation in children with cerebral palsy. *Dev. Med. Child Neurol.* **2021**, *63*, 407–412. [CrossRef] [PubMed]
255. Sharova, O.; Smiyan, O.; Borén, T. Immunological effects of cerebral palsy and rehabilitation exercises in children. *Brain Behav. Immun.-Health* **2021**, *18*, 100365. [CrossRef]
256. Bradley, C.L.; Damiano, D.L. Effects of Dopamine on Motor Recovery and Training in Adults and Children With Nonprogressive Neurological Injuries: A Systematic Review. *Neurorehabil. Neural Repair* **2019**, *33*, 331–344. [CrossRef]
257. Qian, Y.; Forssberg, H.; Heijtz, R.D. Motor Skill Learning Is Associated with Phase-Dependent Modifications in the Striatal cAMP/PKA/DARPP-32 Signaling Pathway in Rodents. *PLoS ONE* **2015**, *10*, e0140974. [CrossRef]
258. Diaz Heijtz, R.; Almeida, R.; Eliasson, A.C.; Forssberg, H. Genetic Variation in the Dopamine System Influences Intervention Outcome in Children with Cerebral Palsy. *eBioMedicine* **2018**, *28*, 162–167. [CrossRef]
259. Worley, G.; Erickson, S.W.; Gustafson, K.E.; Nikolova, Y.S.; Ashley-Koch, A.E.; Belsky, D.W.; Goldstein, R.F.; Page, G.P.; Cotten, C.M. Eunice Kennedy Shriver National Institute of Child Health and Human Development Neonatal Research Network. Genetic variation in dopamine neurotransmission and motor development of infants born extremely-low-birthweight. *Dev. Med. Child Neurol.* **2020**, *62*, 750–757. [CrossRef]
260. Le Pichon, J.-B.; Riordan, S.M.; Watchko, J.; Shapiro, S.M. The Neurological Sequelae of Neonatal Hyperbilirubinemia: Definitions, Diagnosis and Treatment of the Kernicterus Spectrum Disorders (KSDs). *Curr. Pediatr. Rev.* **2017**, *13*, 199–209. [CrossRef] [PubMed]
261. Rose, J.; Vassar, R. Movement disorders due to bilirubin toxicity. *Semin. Fetal. Neonatal Med.* **2015**, *20*, 20–25. [CrossRef] [PubMed]
262. Hamamah, S.; Aghazarian, A.; Nazaryan, A.; Hajnal, A.; Covasa, M. Role of Microbiota-Gut-Brain Axis in Regulating Dopaminergic Signaling. *Biomedicines* **2022**, *10*, 436. [CrossRef]
263. Dicks, L.M.T. Gut Bacteria and Neurotransmitters. *Microorganisms* **2022**, *10*, 1838. [CrossRef]
264. Eisenhofer, G.; Kopin, I.J.; Goldstein, D.S. Catecholamine metabolism: A contemporary view with implications for physiology and medicine. *Pharmacol. Rev.* **2004**, *56*, 331–349. [CrossRef]
265. Bergens, O.; Nilsson, A.; Papaioannou, K.-G.; Kadi, F. Sedentary Patterns and Systemic Inflammation: Sex-Specific Links in Older Adults. *Front. Physiol.* **2021**, *12*, 625950. [CrossRef] [PubMed]
266. Miller, G.E.; Freedland, K.E.; Carney, R.M.; Stetler, C.A.; Banks, W.A. Pathways linking depression, adiposity, and inflammatory markers in healthy young adults. *Brain. Behav. Immun.* **2003**, *17*, 276–285. [CrossRef]
267. Miller, G.E.; Stetler, C.A.; Carney, R.M.; Freedland, K.E.; Banks, W.A. Clinical depression and inflammatory risk markers for coronary heart disease. *Am. J. Cardiol.* **2002**, *90*, 1279–1283. [CrossRef] [PubMed]
268. O'Brien, L.; Hosick, P.A.; John, K.; Stec, D.E.; Hinds, T.D. Biliverdin reductase isozymes in metabolism. *Trends Endocrinol. Metab. TEM* **2015**, *26*, 212–220. [CrossRef]
269. Weaver, L.; Hamoud, A.; Stec, D.E.; Hinds, T.D. Biliverdin reductase and bilirubin in hepatic disease. *Am. J. Physiol.-Gastrointest. Liver Physiol.* **2018**, *314*, G668–G676. [CrossRef]
270. Hinds, T.D.; Gordon, D.M.; Combs, S.D.; Stec, D.E. Bilirubin, a novel endocrine hormone with fat burning properties. *FASEB J.* **2018**, *32*, 605.6. [CrossRef]
271. Li, M.; Kim, D.H.; Tsenovoy, P.L.; Peterson, S.J.; Rezzani, R.; Rodella, L.F.; Aronow, W.S.; Ikehara, S.; Abraham, N.G. Treatment of obese diabetic mice with a heme oxygenase inducer reduces visceral and subcutaneous adiposity, increases adiponectin levels, and improves insulin sensitivity and glucose tolerance. *Diabetes* **2008**, *57*, 1526–1535. [CrossRef]

272. Dekker, D.; Dorresteijn, M.J.; Pijnenburg, M.; Heemskerk, S.; Rasing-Hoogveld, A.; Burger, D.M.; Wagener, F.A.D.T.G.; Smits, P. The bilirubin-increasing drug atazanavir improves endothelial function in patients with type 2 diabetes mellitus. *Arterioscler. Thromb. Vasc. Biol.* **2011**, *31*, 458–463. [[CrossRef](#)] [[PubMed](#)]
273. Dong, H.; Huang, H.; Yun, X.; Kim, D.; Yue, Y.; Wu, H.; Sutter, A.; Chavin, K.D.; Otterbein, L.E.; Adams, D.B.; et al. Bilirubin increases insulin sensitivity in leptin-receptor deficient and diet-induced obese mice through suppression of ER stress and chronic inflammation. *Endocrinology* **2014**, *155*, 818–828. [[CrossRef](#)]
274. Stec, D.E.; Hinds, T.D. Natural Product Heme Oxygenase Inducers as Treatment for Nonalcoholic Fatty Liver Disease. *Int. J. Mol. Sci.* **2020**, *21*, 9493. [[CrossRef](#)]
275. Bigo, C.; Kaeding, J.; El Hussein, D.; Rudkowska, I.; Verreault, M.; Vohl, M.C.; Barbier, O. PPAR α : A Master Regulator of Bilirubin Homeostasis. *PPAR Res.* **2014**, *2014*, 747014. [[CrossRef](#)]
276. Funes, S.C.; Rios, M.; Fernández-Fierro, A.; Covián, C.; Bueno, S.M.; Riedel, C.A.; Mackern-Oberti, J.P.; Kalergis, A.M. Naturally Derived Heme-Oxygenase 1 Inducers and Their Therapeutic Application to Immune-Mediated Diseases. *Front. Immunol.* **2020**, *11*, 1467. [[CrossRef](#)]
277. Diamond, I.; Schmid, R. Experimental bilirubin encephalopathy. The mode of entry of bilirubin-¹⁴C into the central nervous system. *J. Clin. Investig.* **1966**, *45*, 678–689. [[CrossRef](#)]
278. Cui, Y.; Wu, C.; Li, L.; Shi, H.; Li, C.; Yin, S. Toward nanotechnology-enabled application of bilirubin in the treatment and diagnosis of various civilization diseases. *Mater. Today Bio* **2023**, *20*, 100658. [[CrossRef](#)]
279. Yu, M.; Su, D.; Yang, Y.; Qin, L.; Hu, C.; Liu, R.; Zhou, Y.; Yang, C.; Yang, X.; Wang, G.; et al. D-T7 Peptide-Modified PEGylated Bilirubin Nanoparticles Loaded with Cediranib and Paclitaxel for Antiangiogenesis and Chemotherapy of Glioma. *ACS Appl. Mater. Interfaces* **2019**, *11*, 176–186. [[CrossRef](#)]
280. Rahman, A.T.; Shin, J.; Whang, C.-H.; Jung, W.; Yoo, D.; Seo, C.; Cho, B.-K.; Jon, S. Bilirubin Nanomedicine Rescues Intestinal Barrier Destruction and Restores Mucosal Immunity in Colitis. *ACS Nano* **2023**, *17*, 10996–11013. [[CrossRef](#)]
281. Kamothi, D.J.; Kant, V.; Jangir, B.L.; Joshi, V.G.; Ahuja, M.; Kumar, V. Novel preparation of bilirubin-encapsulated pluronic F-127 nanoparticles as a potential biomaterial for wound healing. *Eur. J. Pharmacol.* **2022**, *919*, 174809. [[CrossRef](#)] [[PubMed](#)]
282. Kim, M.J.; Lee, Y.; Jon, S.; Lee, D.Y. PEGylated bilirubin nanoparticle as an anti-oxidative and anti-inflammatory demulcent in pancreatic islet xenotransplantation. *Biomaterials* **2017**, *133*, 242–252. [[CrossRef](#)] [[PubMed](#)]
283. Lee, Y.; Kim, H.; Kang, S.; Lee, J.; Park, J.; Jon, S. Bilirubin Nanoparticles as a Nanomedicine for Anti-inflammation Therapy. *Angew. Chem. Int. Ed.* **2016**, *55*, 7460–7463. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.